

Horticultural News

Volume 91, Number 4
Fall, 2011



Horticultural News

Editors: Winfred P. Cowgill, Jr. & Wesley R. Autio

The New Jersey State Horticultural Society was organized on August 17, 1875 at Geological Hall, Rutgers College, New Brunswick, NJ. It remains the oldest Horticultural organization in New Jersey.



Horticultural News began as the *The New Jersey State Horticultural Society News*, in October of 1920. The Society began "collecting paid membership in order to obtain funds to promote new features of the society and extend the usefulness of the society. The Horticultural Society News was started to be the official society publication." Published M. A. Blake, Professor at Rutgers College was the first president and chair of the publication committee.

Editors served as follows:

MA Blake	1920 - 1947
Norman F Childers	1948 - 1980
Win Cowgill	1981 - 1988
Emily Brown Rosen	1988 - 1990
Linda Butenis Vorsa	1991 - 1995
Jerry Frecon	1995 - 2010

June 2010: *Horticultural News* has moved to an online web-based format. The New Jersey State Horticultural Society has partnered with the University of Massachusetts *Fruit Notes*, Dr. Wesley Autio, Editor. Cowgill and Autio will be the new editors of *Horticultural News* and *Fruit Notes*.

Horticultural News is distributed to growers, extension personnel and researchers and libraries across North America. **Horticultural News** focuses primarily on tree-fruit culture, but addresses small-fruit cultural issues as well. Most reports are from current research at Rutgers University, University of Massachusetts, and other universities.

Horticultural News is published four times per year by the New Jersey State Horticultural Society. It is provided as a benefit to membership in the society. Membership costs \$50 per year. Each one-year subscription begins January 1 and ends December 31. Payments via check must be in United States currency and should be payable to the New Jersey State Horticultural Society. **Horticultural News** Electronic subscriptions are available as benefit of membership in the NJ State Horticultural Society. A hidden link will be mailed immediately after publication to members. Issues will be made freely available on this website six months after publication.

Correspondence should be sent to: New Jersey State Horticultural Society
P.O. Box 116
Clayton, NJ 08312
TEL (856) 307-6450 Ext.1 FAX (856) 307-6476

For advertising opportunities, contact: Meredith Compton njhortsociety@gmail.com

Table of Contents

Preliminary Results with a Vacuum Assisted Harvest System for Apples

James Schupp, Tara Baugher, Edwin Winzeler, Melanie Schupp, and William Messner..... 1

New Cherry Pruning Videos Available Online

Win Cowgill..... 5

Massachusetts Pick-Your-Own Apple Survey

Wesley Autio..... 8

Blueberries - Making a Superb Fruit Even Better!

Stephanie Yao..... 14

Blackberry Variety Selection Opportunities

John Clark..... 21

New Jersey News..... 24

Cover: Primocane-fruiting blackberries at the UMass Cold Spring Orchard Research & Education Center, Belchertown, Massachusetts. Photo by Wesley R. Autio.

Preliminary Results with a Vacuum Assisted Harvest System for Apples

James Schupp, Tara Baugher, Edwin Winzeler, and Melanie Schupp
College of Agriculture, Penn State University

William Messner
Department of Mechanical Engineering, Carnegie Mellon University

The harvest of specialty crops for the fresh market is labor intensive, and attempts at automation have been less successful than with field crops. Apple harvest is particularly difficult to automate because fruit suffer bruise damage easily. Nevertheless, market, social, and political forces have converged to make mechanical augmentation of harvest essential for the survival of the specialty crop industry in the U.S.

The challenges are enormous, as the constraints on candidate technologies include high performance, low cost, robustness, simplicity, and ease of repair. The opportunities and rewards, on the other hand, are commensurately great. Merely addressing these challenges is already inspiring a new generation of engineers and students to think creatively about problems in agriculture and related fields and to bring engineers and growers together (Kliethermes et al., 2010; Leslie et al., 2008; http://www.cascrop.com/index.php?option=com_content&view=article&id=1521&Itemid=666). Successful development technologies could reinvigorate the specialty crop industry, make it competitive in international markets, and employ segments of the population that have largely been excluded from the labor pool due to physical constraints.

The total value of U.S. specialty crops—\$49 billion in sales—now exceeds the combined value of the five major program crops—\$45.8 billion in sales (Schmoldt, 2007). However, despite the specialty crop industry's major contribution to the U.S. economy and the finding that “a secure domestic food supply is a national security imperative,” U.S. specialty crop producers remain vulnerable to the real possibility of being eliminated within the next ten years (Schmoldt, 2007). This crisis stems in large part from dependency on a large seasonal workforce, coupled with increasing labor costs and decreasing availability of agricultural employees. In a

socioeconomic technology adoption survey of growers conducted by members of our research team, harvesting was among the highest rated areas of need for advanced technologies to improve precision and efficiency in tree fruit production (Ellis et al., 2010).

Prior Approaches to Addressing Harvest Labor Inputs

Mechanical harvesting machines that utilized mass removal techniques were widely tested in the U.S. in the 1970s and 1980s. The machines were unsuccessful in harvesting fruit for the fresh market due to excessive fruit damage caused during fruit detachment, contact with limbs or other fruit while falling through a three-dimensional tree canopy, and bulk collection procedures (Peterson, 2005b).

Mechanical engineering efforts for specialty crops declined in the 1990s, and the focus shifted to the development of labor platforms for use with planar tree architectures. Fruit were still picked and placed in the bin by hand, but harvest efficiency was increased and fruit quality was similar to that which was conventionally harvested (Baugher et al., 2009a; Schupp et al., 2007). In the late 1990s, engineers began looking at automated bin filling technologies, but early designs resulted in excessive bruising of fruit (Peterson, 2005a). The complex fruit handling and equipment/operator interface was a major obstacle to developing semi-automated harvest systems.

Significant progress has been made on robotic harvest. However, insufficient fruit recovery and difficulties in developing both an end effector and a vision system that performs equal to the human hand and human visual system avert commercialization in the near future (Bulanon and Kataoka, 2010; Sarig, 1993).




Figure 1. A. Components of vacuum-assisted harvest system. B. Elephant-ear distributor in motion.

Current Research with Commercial Partner

In Fall 2011, with support from the Specialty Crop Research Initiative Project, “Comprehensive Automation for Specialty Crops” (Singh et al., 2011), our harvest team began working with a commercialization partner, DBR Conveyor Concepts, on a vacuum tube transport system and automated bin filler that could be retrofitted to existing grower equipment. Figure 1 shows the first prototype with the major components labeled.

Suction is provided by a pair of vacuum pumps driven by an internal combustion engine. The pumps and engine are in an enclosure mounted on the work platform. The vacuum pumps exhaust through a vent on the top of the enclosure. The exhaust pipe of the internal combustion engine is also at the top of the enclosure. Vacuum return hoses (green) lead from the pumps to the proprietary deceleration mechanism, which is the key innovation of this system. The vacuum pumps lower the internal pressure of the deceleration mechanism enclosure below ambient, causing air to flow through the vacuum hoses (black). Pickers place apples into the inlet to the vacuum hoses opposite the deceleration mechanism. The vacuum hoses are padded to prevent bruising of the fruit.

When a picker (partially hidden by the tree) places an apple into the vacuum hose, air flow into the hose is obstructed, leading to a differential pressure across the apple. The unequal pressure forces the apple through the hose. When it reaches the enclosure of the deceleration mechanism, its momentum propels into the deceleration mechanism, which has two functions. The first function is slowing down the apple and dropping it onto the flexible-flap bin filling mechanism (occluded by the bin in Figure 1A; shown in motion in Figure 1B). The second is to provide an airtight seal between the portions of the transport mechanism that are held below ambient pressure and the exit port of the deceleration mechanism of the enclosure.

Materials & Methods

Initial trials were conducted to assess fruit bruising at various stages in the augmented harvest system—(1) prior to entry into the vacuum tube (the control treatment), (2) after the vacuum tube and decelerator and before the elephant ears, (3) after the elephant ears but before the bin, and (4) after transport through the entire system. Bruising and corresponding USDA fruit grades

were assessed as described in Kliethermes et al., 2010. Five replicates of either 15 or 20 fruit were randomly subjected to each of the treatments. The studies were first conducted on ‘Honeycrisp’ and ‘Daybreak Fuji.’ Based on the bruise findings, modifications were made to the harvest system to further prevent bruising, and a final study was conducted on ‘Golden Delicious,’ which is highly bruise-susceptible.

Commercial-scale efficiency trials were conducted on ‘Golden Delicious,’ ‘York,’ and ‘Pink Lady’ to assess labor productivity and fruit quality in apple orchard plots harvested with the vacuum assist system and a work platform compared to hand harvest and ladders. The same workers performed both treatments within a trial. The experimental design was randomized complete block with four multiple-tree replicates. Harvest times were compared for each treatment, and bruise evaluations were conducted on 100 fruit per treatment. Data from all trials were subjected to an analysis of variance and means were separated using Fisher’s protected least significant difference test.

Results & Discussion

The initial two trials to assess fruit damage on apples collected at various stages in the augmented harvest system revealed that changes should be made to the elephant ears to prevent bruising (Table 1). With ‘Honeycrisp,’ bruise volume in fruit collected from the elephant ears was higher than control fruit collected prior to the vacuum tube, although the effect on fruit grade was insignificant. In the trials with ‘Daybreak Fuji,’ bruise volume in fruit collected from the bin was higher than fruit collected prior to the vacuum tube, and the portion of fruit that graded U.S. Extra Fancy was reduced from 99 percent to 92 percent. Machine modifications to further reduce fruit damage eliminated bruising in the final trial with ‘Golden Delicious,’ and fruit graded almost 100 percent U.S. Extra Fancy.

Commercial-scale investigations on ‘Golden Delicious,’ ‘York,’ and ‘Pink Lady’ demonstrated increases in efficiency per acre of 10 to 49% (Table 2). The quality of machine-harvested fruit was equal to hand harvested fruit in the ‘Golden Delicious’ and ‘Pink Lady’ trials and was better than hand harvested fruit in the ‘York’ trial (Table 2). The cost/benefit beyond hand harvest was \$245 to \$517 per acre (Baughner et al., 2011; data not shown; www.cascrop.com).

From an engineering perspective, the vacuum as-

Table 1. Bruise volume measured on apples collected at various stages in augmented harvest system and corresponding effects on fruit graded USDA Extra Fancy.

Cultivar	Location of Sample	Bruise volume (mm ³)	U.S. Extra Fancy ^z (%)
Honeycrisp	Before vacuum tube (control)	2.0 b	
	After vacuum and decelerator	12.8 ab	96.0 a ^x
	After elephant ears	20.1 a	94.7 a
	From bin	8.3 ab	96.0 a
Daybreak Fuji	Before vacuum tube (control)	0.0 b	
	After vacuum and decelerator	13.5 b	98.7 a
	After elephant ears	14.7 ab	98.7 a
	From bin	34.6 a	92.0 b
Golden Delicious ^y	Before vacuum tube (control)	2.5 a	
	After vacuum and decelerator	7.9 a	99.9 a
	After elephant ears	10.4 a	99.8 a
	From bin	17.0 a	99.9 a

^z Mean separation within columns and cultivars by Fisher's protected least significant difference at P=0.05 (Five replicates in each trial; n=20, Golden Delicious; n=15, Honeycrisp, Daybreak Fuji).

^y Vacuum tubes, decelerators, and elephant ears modified to further reduce bruising prior to Golden Delicious trial conducted on October 11, 2010.

bruising. The deceleration mechanism solves the important problem of isolating the vacuum from ambient pressure while providing a soft ejection for fruit. The modular design of the entire system makes it attractive to use with standard orchard equipment such as platforms and bin trailers.

At the same time there is room for further improvement. The biggest problems are that the vacuum pumps and the internal combustion engine are quite noisy, and working near the engine is hot. Another problem is that there are only two vacuum hoses. More hoses will be needed to allow more pickers to work at the same time to make the system cost effective. Our commercial partner is addressing these issues with their next prototype which will employ a single, larger but slower moving (and therefore quieter) vacuum pump driven by a hydraulic motor, with hydraulic power provided by the tractor towing the

sisted transport system addresses a number of design challenges well. The entire system is simple, uses readily available materials and parts, and thus it is easy to maintain or repair. The vacuum tube and deceleration mechanism effectively move apples from the picker to the bin filling device with minimal

Table 2. Labor efficiency and fruit quality in apple orchard plots harvested with vacuum assist system and platform compared to hand harvest and ladders.

Cultivar	Harvest System	Harvest Time ^z (h/acre/person)	Efficiency (% increase)	Fruit downgraded ^y (%)
Golden Delicious	Vacuum assist	33.35 b ^x	9.8	11.1 a
	Hand	36.98 a	--	15.6 a
York	Vacuum assist	24.69 b	49.2	6.0 b
	Hand	48.60 a	--	10.6 a
Pink Lady	Vacuum assist	37.47 b	19.4	5.3 a
	Hand	44.74 a	--	8.1 a

^z Includes harvest of lower portion of trees by hand.

^y Percentage of fruit downgraded determined from bruise evaluations conducted on 100 fruit per treatment from each of four replicates.

^x Randomized complete block. Mean separation within columns by Fisher's protected least significant difference at P=0.05.

system. The new system will feature four vacuum hoses and deceleration devices. These improvements may be enough to make the vacuum assisted harvester not only viable technologically, but also economically profitable for growers.

Acknowledgements

The authors acknowledge the valuable contributions of Phil Brown, Mike Rasch, Chuck Dietrich, Terry Salada, Freeman Showers, and Eric Anderson.



Literature Cited

- Baughner, T. A., K. Lewis, J. Schupp, K. Lesser, M. Harsh, C. Seavert, T. Auvil. 2009a. Mobile platforms increase orchard management efficiency and profitability. *ACTA Horticulturae* 824:361-364.
- Baughner, T. A., J. Schupp, E. Winzeler, W. Messner, M. Bergerman. 2011. Mechanically Assisted Harvest of Apples. Proc. American Society of Agricultural and Biological Engineering Annual Convention. Abstract.
- Bulanon, D.M. and T. Kataoka. . 2010. Fruit detection system and an end effector for robotic harvesting of Fuji apples. *Agricultural Engineering International: CIGR Journal* 12(1):203-210.
- Ellis, K., Tara Auxt Baughner, K. Lewis. 2010. Use of survey instruments to assess technology adoption for tree fruit production. *HortTechnology* 20:1043-1048.
- Kliethermes, B., A. Leslie, R. Rohrbach, J. Koan, S. Wolford, M. Glenn, K. Lewis, T. Baughner, and W. Messner. 2010. Novel approaches to passive bin filling for apples. ASABE paper 1009399. 16 pp.
- Leslie, A., R. Rohrbach, J. Koan, B. Kliethermes, W. Messner, T. Baughner, and P. Heinemann. 2008. Performance of energy absorbing materials for passive bulk bin filling. *Proc. Cumberland-Shenandoah Fruit Workers Conference*. Winchester, VA. 9 pp.
- Peterson, D. L. 2005a. Development of a harvest aid for narrow-inclined-trellised tree-fruit canopies. *Applied Engineering in Agriculture* 21(5):803-806.
- Peterson, D. L. 2005b. Harvest mechanization progress and prospects for fresh market quality deciduous tree fruits. *HortTechnology* 15(1):72-75.
- Sarig, Y. 1993. Robotics of fruit harvesting: A state-of-the-art review. *Journal of Agricultural Engineering Research*. 54(4):265-280.
- Schmoldt, D. 2007. The interplay of biology and engineering for smarter applications. *Biological Engineering* 1(2):111-125.
- Schupp, J. R., T. A. Baughner, R. M. Harsh, K.M. Lesser, and B.D. Wenk. 2007. Mobile platforms increase orchard labor efficiency. *HortScience* 42:4. (July) Abstract.
- Singh, S., M. Bergerman, G. Hoheisel, K. Lewis, T. Baughner. 2011. Comprehensive automation for specialty crops (CASC) – developing a more sustainable and profitable U.S. specialty crop industry. <http://www.cascrop.com/>.

New Cherry Pruning Videos Available Online

<http://giselacherry.com/>

<http://www.youtube.com/giselacherry>



PLAN AHEAD - DON'T BE LEFT OUT!

Are You Prepared for Your Harvest of Tomorrow?

PLAN AHEAD

2013 BUDDING CONTRACTS DUE JUNE 1st!

Order dwarf cherry trees NOW for 2011-2012

Use our **knowledge** and **experience** to secure
your tree needs. **All at no cost to you.**



SUMMIT TREE SALES

REPRESENTING THE COUNTRY'S FINEST NURSERIES

800.424.2765 / 269.674.8866 / Fax 269.674.3758

summit@summittreesales.com



Wally Heuser



Wanda Heuser Gale



Jan Melvin



Helen Classen



Katie Schuld



Dawn Melvin

s u m m i t t r e e s a l e s . c o m

Targit Sales

"Specializing in Farm Market Packaging"

1-800-526-9224

Biodegradable

PLASTIC BAGS



Targit Sales is proud to introduce our line of Biodegradable plastic bags. Our bags use a proprietary additive that renders the plastic ultimately biodegradable.



We are offering the Biodegradable bag program with all of our "Custom Imprinted" plastic bags. Our Biodegradable bags have the "EPI" (the additive company) logo on the bag as proof of its biodegradable properties.

Advertise your business along with a "tag line" to express your company's efforts to keep your community and planet green.

Targit Sales can fully customize your bags to your specifications. We can make any size, strength, and style of bag to fit your needs. We can assist you with your design, setup, and artwork.

COMMON SIZES OF BAGS:

- 13 x 10 x 24 (Great for pies, cakes and corn)
- 12 x 8 x 24 (Great for your store front)
- 8 x 5 x 21 (Perfect size for tailgate markets)

BIODEGRADABLE PLASTIC BAG BENEFITS!

- * Superior Strength!
- * Less expensive to make and ship versus paper bags!
- * Superior imprinting capability and quality...(looks better)!
- * Easier and more affordable to store (inventory)!
- * Flexibility! (customize the size, thickness, color and style to your specifications)
- * Reusability!
- * Biodegradable..breaks down quickly!

MOST COMMON QUESTIONS asked about biodegradable bags.

Will the bags begin to break down while being stored? No, the bags will retain its strength and integrity while properly stored. The breakdown only begins after it is directly exposed to certain environmental conditions such as: soil/landfill, heat/moisture, and sunlight.

Is this process more expensive? Not really, the cost increase is only a few percentage points higher. You will more than make this up with the goodwill you will be conveying to your community and customers.

Does the bag lose its strength or feel...does it look different? No, it is almost impossible to tell the difference between the Biodegradable bags and the traditional plastic bags. This is why we recommend putting a tag line on your printed biodegradable bags.

Can these bags be recycled and/or reused? Yes, these bags can be recycled and reused.

For more information or to order contact
Targit Sales
1-800-526-9224 • www.targitsales.com

Targit
sales associates
"Specialty Food Packaging"

Massachusetts Pick-Your-Own Apple Survey

Wesley R. Autio


Department of Plant, Soil, & Insect Sciences, University of Massachusetts

Orchardists have long recognized that a good awareness of what others are using for orchard practices is an excellent guide for the development of their own practices. This recognition extends to all aspects of the orchard operation, from tree planting to sales. To get a better understanding of how Massachusetts apple growers work with pick-your-own sales, at the December 2011 Massachusetts Fruit Growers' Association Directors Meeting, Alex Dows proposed that we survey Massachusetts apple growers. The Directors enthusiastically supported the idea, and a draft survey was distributed among the Directors in late December and early January. After some modification, the final survey was developed.

To aid in the process of distribution, SurveyMonkey.com was used to house the survey, distribute the survey, and collect the data. This tool allowed us to track responses and to follow up those who had not responded with additional requests. In all, survey requests

Table 1. Pick-your-own survey distribution and response rate.


Surveys distributed (no. of orchards)	Survey responses (no. of orchards)	Survey responses (% of orchards)
110	83	75%
Massachusetts total acreage	Survey responses (acres)	Survey responses (% of Mass. acreage)
4287	1798	42%



were distributed to 110 individuals (members of the Massachusetts Fruit Growers' Association and subscribers to the UMass *Healthy Fruit* newsletter). Of those distributed, 83 responses (75%) were received (Table 1). Within the responses, 1798 acres of apple production were represented (about 42% of the Massachusetts total, Table 1).

Among responses, the average farm had 28 acres of apples. Sixty-four percent of farms used pick-your-own (PYO) sales, and on average more than half of the acreage of those farms was devoted to PYO (Figure 1).

About 30% of the PYO acreage is in dwarf trees, 53% in semi-dwarf, and 17% in standards (Figure 2). Most farms provide access




mazes are part of the entertainment at just under 10%. Several other forms of entertainment are used, including walking trails, cider pressing, pony rides, rock climbing walls, kids games, BBQs, snack bars, farm stores, picnic areas, and speed dating.



Advertising is a critical part of nearly all PYO orchards (Figure 5). Everyone uses word-of-mouth advertising. Seventy-five percent use newspaper, 50% use email lists, 25% use radio ads, 15% use direct mailings, and 11% use television ads. Sixty-five percent of farms use other forms of advertising, such as the internet (43% use web-based tools), road signs, PYO guides, flyers, and brochures. Among the various techniques, the greatest portion of the advertising dollar (50%) is spent on newspaper ads (Figure 6).



for handicapped people, and only 17% provide some form of access aid, such as ladders (Figure 3).

About 30% is spent on email lists, radio, direct mailings,

School tours are provided by 80% of PYO farms (Figure 4). On average, farms charge \$5.17 per child for school tours. Many farms utilize other forms of entertainment to supplement PYO visits (Figure 4). Hayrides are made available to customers at 66% of PYO farms. Animals are an attraction at 36%. Live music is used at more than 20% of farms, and various types of







and television together.

Among PYO orchards, 87% sell fruit by the container, with an average charge of \$13.05 per peck. Of the small number selling by the pound, the average

Table 2. Pros and cons (from the farmer’s perspective) associated with pick-your-own sales as listed by Massachusetts PYO Survey respondents in 2011.

Pros	Cons
Beautiful views	Crowds
Easy access	Waste
Family friendly	The public
Entertainment	Theft
Repeat customers	Weather dependent
More time for sales	Vandalism
Freedom of choice	Risk
Just a "real orchard"	Parking
Happy customers	
Attracts customers	
Keeps farm viable	
Dog friendly	
Good return	
Less physical labor	
Great exposure	
Extends season	
Immediate cash flow	




Figure 8. Extent of entry fee usage and the timing of payment in Massachusetts PYO orchards.

charge is \$1.42 per pound. Entry fees are charged by fewer than 10% of all PYO orchards, and 68% charge the customer for fruit prior to picking.


The survey included three very open-ended questions. The first asked farmers to list pros and cons associated with the PYO aspects of their farm (Table 2). The pros were all as you might expect, and included “happy customers,” “great attraction,” “less physical labor,” etc. The cons also were as you might expect, and include the problems associated with crowds, theft, parking, and the weather.

A second open-ended question asked farmers what changes they were anticipating in the near future (Table 3). PYO orchards are considering more parking, more varieties, added entertainment, and better orchard management. One noted that their future plans primarily include retirement!

The third open question asked farmers to list their most serious concerns for the future of their PYO operation (Table 3). The three most serious concerns were regulation, food safety, and this survey encouraging

Table 3. Planned changes and concerns (from the farmer’s perspective) associated with pick-your-own sales as listed by Massachusetts PYO Survey respondents in 2011.

Changes	Concerns
Better crowd control	Regulation
More parking	Food safety
Improvement of animal area	This survey
Enhanced handicap access	
New varieties	
Donut machine	
Increase price	
More dwarf trees	
One-price bags	
Improved stand	
Better signage	
Loyalty discounts	
Email lists	
Hay rides	
U-sort apples	
Expand plantings available	
More value-added products	
Better orchard management	
Retirement	



more competition.

The last question on the survey asks growers to rate their level of satisfaction with PYO on their farm (Figure 9). Overwhelmingly, orchards were satisfied or very satisfied with PYO. Seventeen percent were neutral, and 8% were dissatisfied with PYO.

We hope that this survey helps guide the future develop of PYO as an apple sales technique.



Maximize Your Fertilizer Efficiency and Crop Quality.

System-Cal

System-Zinc

System-Mag

System-Manganese

System-K



Agro-K, the premier name in manufacturing high quality foliar nutrients world-wide for over 30 years has partnered with CPS to provide Northeast fruit and vegetable growers with the nutrient tools they need maximize crop quality – size, firmness, storage life, and more. Agro-K produces a full line of quality foliar nutrient products including the **System™ line** of phosphite-based micronutrients (including calcium, zinc, magnesium, manganese and potassium), to help growers improve their ground fertilizer efficiency, overall plant health and crop quality. Agro-K also offers a complete line of OMRI approved nutrients for use in organic farming.

For more information contact your local CPS crop specialist or Agro-K's Northeast Regional Mgr., Jeff McClellan at 814-574-5663 or jeff@agro-k.com.

AGRO-K CORPORATION
8030 Main Street, NE • Minneapolis, MN 55432
800-328-2418 • www.agro-k.com



Blueberries - Making a Superb Fruit Even Better!

Stephanie Yao

Formerly with Agriculture Research Service, USDA



Blueberries are popular and versatile—you can put them in or on almost anything. But the berry would not be where it is today without the efforts ARS researchers. Today, ARS scientists are busy solving growers' problems with blueberry disease, firmness, splitting, and cold tolerance.

When U.S. Department of Agriculture botanist Frederick Coville started the world's first successful blueberry breeding program, did he envision it would grow into the multi-million dollar industry it is today? Maybe. But a century later, thanks to dedication by

Coville, collaborator Elizabeth White, and other USDA and university scientists, blueberries are the second most popular berry consumed in the United States.

A member of the genus *Vaccinium*, blueberries are related to many commercially important and popular fruit species, like cranberry, lingonberry, and huckleberry. Blueberries are mainly native to North America and are lauded for their health benefits.

Coville began researching blueberries in 1906, when he started a series of experiments to learn fundamental facts about them, thinking they might be suitable for cultivation. Coville found that blueberries and many other plants require acid soils to grow, a fact not known to horticulturists prior to his experiments.

After a few years of study, Coville published in 1910 the first bulletin outlining how to successfully grow blueberries from seed to fruit. White, whose family at that time had a successful cranberry farm in New Jersey, helped Coville acquire some of the best wild blueberry plants to use as parents in his breeding experiments.

In 1911, Coville made the first cross of



Plant geneticist Mark Ehlenfeldt (left) and plant pathologist James Polashock examine blueberry plants and collect data on mummy berry fruit infection to evaluate resistance.

wild blueberry germplasm that eventually led to the release of several blueberry cultivars—ancestors of cultivars currently grown throughout the world—marking the beginning of USDA’S current breeding program.

Throughout the years, notable Agricultural Research Service blueberry breeders George Darrow, Donald Scott, and Arlen Draper have made significant contributions to the advancement of blueberries. Today, 100 years after Coville made his first successful cross, ARS researchers throughout the country continue that longstanding goal of improving blueberries so consumers can enjoy them for many more centuries to come.

Mitigating Mummy Berry Blight & Fruit Rot


Geneticist Mark Ehlenfeldt and plant pathologist James Polashock are researching mummies—mummified blueberries, that is, which got that way because of a disease. The scientists are with the Genetic Improvement of Fruits and Vegetables Laboratory in Beltsville, Maryland, and are stationed at the Philip E. Marucci Center for Blueberry and Cranberry Research and Extension in Chatsworth, New Jersey. One of ARS’s flagship locations for blueberry research, Chatsworth houses the largest collection of potted and in-ground blueberry cultivars in the world.

In addition to releasing improved blueberry varieties, the researchers focus on screening for disease resistance, and mummy berry is one of the most important blueberry diseases in North America.

“Mummy berry is caused by the fungus *Monilinia vaccinii-corymbosi*,” says Polashock. “It occurs almost everywhere blueberries are grown and affects all cultivated species, including highbush, lowbush, rabbiteye, and some wild species.”

Mummy berry disease is unique because it occurs in two distinct phases. During the blighting phase, small, cup-shaped structures bearing fungal spores sprout from mummified berries concealed in leaf litter on the ground. Wind spreads the spores to blueberry plants, infecting the newly emerging shoots and leaves. A second phase of spores, produced on blighted tissue, is carried by bees to the flowers, beginning the fruit-rotting stage. During this phase, the fungus fills the inside of the blueberry as it grows and causes it to shrink, shrivel, and turn whitish—hence the mummy reference. The mummified fruit drops to the ground and overwinters, waiting to begin the process again in the spring.

In an effort to mitigate this disease, Ehlenfeldt, Polashock, plant pathologist Allan Stretch (now retired),



James Polashock screens blueberry tissue cultures for plantlets that have transformed, or changed, their genetic makeup. These plantlets are easy to identify because they express a green fluorescent protein and glow under UV light in the procedure being used. In these transformed plantlets, the genes that respond to the fungus that causes mummy berry are likely to provide clues to resistance to the disease.

and statistician Matthew Kramer undertook two long-term, simultaneous studies examining cultivar response. The first study, published in the scientific journal *HortScience*, sought to predict cultivar resistance and susceptibility to both phases of the disease. The scientists examined more than 90 blueberry cultivars over 9 to 12 years.

“We found that disease response had significant and large genotype-by-environment interactions,” explains Ehlenfeldt. “This means that the 2-3 years of data typically used for publication aren’t enough to reliably estimate disease resistance. Breeders should be evaluating resistance for 8 years to get a good estimate of cultivar response to this disease.” The researchers found an important predictor of blighting to be either the average amount of precipitation at the end of January or rain frequency at the end of March. The average high temperature in late February was predictive for the fruit-infection phase.

Despite predictions of needing 8 years to estimate disease resistance, a second study, also published in *HortScience* analyzed data from 125 cultivars tested for 2-6 years for resistance to the blighting phase and 110 cultivars tested for 2-5 years for resistance to the fruit-infection stage. Using innovative statistics developed by Kramer, the researchers were able to rank resistances among the wide range of cultivars. “For breeding, one

often needs only to know which cultivars are the most resistant on a relative basis,” says Ehlenfeldt. They found several cultivars, such as Brunswick and Bluejay, to be resistant to both phases of mummy berry infection.

“Ultimately, documentation of resistance to each phase will help growers select which cultivars to plant,” says Ehlenfeldt. “This will also help breeders develop strategies to produce cultivars with superior resistance.”

Preventing Fruit Splitting

The Thad Cochran Southern Horticultural Laboratory in Poplarville, Mississippi, joined ARS’s blueberry research program in the 1970s. Led by horticulturist James Spiers (now retired), the program was started after the region’s tung oil industry collapsed because of competition from imported petroleum and a devastating blow from Hurricane Camille in 1969. “Rabbiteye blueberries are native to the Southeast,” says Spiers. “ARS has also introduced a southern highbush blueberry to the region. Combined,



Horticulturist Donna Marshall measures blueberry firmness to determine the correlation between fruit firmness and susceptibility to fruit splitting.

the two blueberry species have proven to be a viable specialty crop for this area.”

So far, Poplarville scientists have released 15 cultivars for growers in the Southeast. But that’s not all they do. The researchers also focus on solving problems growers face, such as rain-induced fruit splitting.

“Splitting and cracking occur in southern highbush and rabbiteye blueberries if they receive preharvest rainfall when fully ripe or approaching ripeness,” explains horticulturist Donna Marshall. She works with Spiers, geneticist Stephen Stringer, and University of Southern Mississippi associate professor Kenneth Curry on this problem. “Researchers have studied rain-induced splitting in cherries, grapes, and tomatoes, but it hasn’t been explored in blueberries.”

Splitting can be mild, in the form of a shallow crack in the skin, to severe, such as deep wounds that penetrate the pulp. But regardless of severity, all splitting renders the fruit unmarketable. Growers in Mississippi and Louisiana have reported as much as 20 percent crop loss on highly susceptible cultivars. That amounts to losses of \$300 to \$500



ARS researchers in Corvallis, Oregon, are developing and improving blueberries for the Pacific Northwest. Shown here are Elliott blueberry plants in full bloom.



Close-up of blueberry flowers.

“Through our studies, we’ve shown that splitting is a cultivar-specific problem,” says Marshall. “But there are still questions, such as what is going on at the cellular level that allows a cultivar to stay intact? With further research, we hope to find the answer.”

Generating Genomic Tools for Blueberry Improvement

Geneticists Chad Finn, with the ARS Horticultural Crops Research Unit, and Nahla Bassil, with the ARS National Clonal Germplasm Repository—both in Corvallis, Oregon—are developing

per acre.

The researchers examined several aspects of fruit splitting in three studies published in *HortScience*. In the first study, published in 2007, the researchers developed a laboratory method to model rain-related splitting in blueberries. Many breeders throughout the country are using this method to more vigorously screen cultivars and selections for splitting susceptibility. The results from field and laboratory tests showed that the rabbiteye cultivar Premier had the lowest incidence of splitting, while widely grown cultivar Tifblue exhibited a high incidence of splitting.

Marshall and colleagues also investigated the correlation between splitting susceptibility and fruit firmness. Laboratory and field tests proved that, in general, firmer fruit has a higher tendency to split. But one selection, named “MS614,” exhibited extreme firmness and splitting resistance. The results, published in 2008, suggest that breeders who select for firmness may inadvertently also be selecting for splitting. But the laboratory screening method Marshall and colleagues created has helped remedy this problem.

The most recent study, published in 2009, evaluated water-uptake thresholds in split-resistant Premier and split-susceptible Tifblue fruit at all stages of development. The researchers harvested and weighed the fruit, then soaked it in distilled water at room temperature for 24 hours. They found that Premier absorbs more water than Tifblue yet remains intact and experiences minimal splitting.



Fruit cluster of Draper, a cultivar released by Michigan State University and named in honor of Arlen Draper, a long-time blueberry breeder with ARS in Beltsville, Maryland.

and improving blueberries for the Pacific Northwest. Although Corvallis is the most recent ARS location to conduct blueberry breeding, Finn and Bassil are playing an important role in a nationwide, multi-institutional project aimed at developing genomic tools to help improve blueberries.

Funded by the Specialty Crops Research Initiative, the project is led by fellow ARS geneticist Jeannie Rowland in Beltsville, Maryland, and involves several university and international collaborators. Finn and Bassil are working with Michigan State University professor James Hancock in developing a genetic map for highbush blueberry.

“We are currently testing plants made from a cross between the northern highbush cultivar Draper and the southern highbush cultivar Jewel at various locations

across the country where blueberry is grown,” says Finn. “Our task is to compare the performance of each plant in the field. For the next couple of seasons, we will evaluate the plants for chilling requirement, cold tolerance, and fruit-quality traits.”

In the lab, Bassil is processing leaf samples to extract DNA and genotype the plants. The researchers will then merge the field and lab data to determine whether genetic markers that predict a plant’s performance can be identified. Bassil is also helping to develop genetic markers and following them through mapping populations and wild blueberry populations for genetic diversity studies.

The new tools, once available, should make blueberry breeding and cultivar development far more efficient.

Source: Agricultural Research -- May-June 2011
(<http://www.ars.usda.gov/is/AR/>)

Specialized. Diverse. Innovative.

At Adams County Nursery we specialize in commercial fruit tree nursery stock for a diverse, eastern market using the latest in technology and innovation.



Adams County Nursery, Inc.
(800) 377-3106 • (717) 677-4124 fax
www.acnursery.com • email: acn@acnursery.com

Orchard Equipment and Tools for Every Grower



REARS
REARS Pulblast

Jacto

CARRARO SPRAY
by CARRARO

ATVM
ATV

MS
MS 3-PT Hitch Boom Sprayer

OESCO, INC.

Parts & Service for all we sell!
P.O. Box 540, Rte. 116 • Conway, MA 01341
800-634-5557 • 413-369-4335
info@oescoinc.com

Order online at www.oescoinc.com

Since 1932

The Best Berry Plants



80 YEARS

- Strawberries, raspberries, blueberries, blackberries, asparagus and more!
- Where the pros go for plans and plants.
- Call for a free catalog and plasticulture guide!

41 River Road, South Deerfield, Massachusetts 01373

NOURSE

www.noursefarms.com 413.665.2658

RIGHT...FROM THE START

Start your orchard RIGHT with Willow Drive Nursery's Proven Performers™. Call for a complete and updated inventory.

Calling us today can help you tomorrow!



APPLES

Braeburn

Mariri Red™ (Eve®)
Kumeu Crimson®

Fuji

Aztec Fuji® (D'Y 2 Variety)
Rising Sun®
Morning Mist™
Early Fuji

Gala

Brookfield® Gala
Buckeye® Gala
Ultima Gala®

Other

Cameo® brand
Granny Smith
Honeycrisp™

It® Red Delicious
Morren's® Jonagored
Supra™
Ruby Jon®
RubyMac®
Smoothee® Golden
Spartan



CHERRIES

Attika®

Benton™

Bing

BlackGold™

BurgundyPearl™

Chelan™

Early Robin®

EbonyPearl™

Lapins

Montmorency

Rainier

Regina

Rynbrandt

Schmidt

Selah™

Skeena™

Sweetheart™

Tieton™

Ulster

WhiteGold™



PEARS

Bartlett

Columbia Red Anjou™

Concorde™

Comice

D'Anjou

Forelle

Golden Russet Bosc®

Red Clapp's Favorite



PEACHES

Stellar Series

Allstar®

Autumnstar®

Brightstar™

Coralstar®

Earlstar™

Flamin' Fury

PF 8 Ball

PF Lucky 13

PF 19-007

PF 25

PF 35-007



WILLOW DRIVE NURSERY, INC.

1-888-54-TREES

Ephrata, Washington · 1-888-548-7337 · Web Site: www.willowdrive.com

Blackberry Variety Selection Opportunities

John R. Clark

Department of Horticulture, University of Arkansas

Blackberries offer another small fruit option for eastern US growers for a range of markets. The major issues for Pennsylvania and Mid-Atlantic growers when considering blackberry varieties include winter hardiness, maturity date, quality, yield, berry size, and overall adaptation. I will share some information in general about blackberries, some market and production changes in the US, and finally some variety thoughts for Mid-Atlantic growers.

Blackberry and raspberry plants are rather unique in the fruit world in that they have a perennial root system but have biennial canes. This means the canes live two years and then die. The two cane types are primocanes, or first-year canes, and floricanes, which are second-year canes. In red raspberry, primocane- and florican-fruiting varieties exist and both have substantial commercial value. In blackberries, florican-fruiting has been the basis of all blackberry production and commercial primocane-fruiting varieties did not exist prior to the release of Prime-Jim®, Prime-Jan®, and Prime-Ark® 45 by the University of Arkansas. Primocane fruiting offers the opportunity for late summer and fall production, to complement summer production of blackberries. However, there is much to be done in the improvement of primocane blackberries and all the answers are not in place yet for Mid-Atlantic growers.

Aspects of Blackberry Production History

A survey of eastern U.S. (east of the Rocky Mountains) production in 1990 indicated blackberries were marketed in the following ways: 62% pick-your-own, 36% pre-picked fresh market (mainly on-farm or local fruit stand sales), and 2% processed. The survey results did not indicate that any production was for shipping to distant markets or grocery stores. In this survey, production area increased 56% from 1980 to 1990, with a further projected 66% increase in production area from 1990 to 2000. Again, this increase was envisioned to be largely marketed locally. In the early 1990s, blackberries were not found on grocery store shelves across the U.S. (some were present on the west coast), due

mainly to the lack of postharvest handling capability of varieties released prior to that time, but also because blackberries had never made much headway into the competition for commercial grocery shelf space.

Some of the new varieties beginning to be planted in the early 1990s were found to have fruit firmness adequate for shipping. Chester Thornless displayed a good level of firmness and shelf life, and in the world picture became a major shipping berry later in the decade. The Arkansas-released Navaho was found to have excellent shelf life. Subsequent releases from Arkansas included 'Arapaho' and 'Apache', each of which had fruit capable of withstanding shipping. These varieties contributed to a major shift in the production outlook for shipping of blackberries from that of a local-marketed crop to one shipped for retail marketing.

A major development occurred during the 1990s: the shipping of blackberries to the U.S. from Chile and Guatemala. Soon thereafter, the development of production technology in Central Mexico increased availability of eastern US-developed blackberries. Mexican production is centered in the highlands of the state of Michoacan and Jalisco, and utilizes a number of techniques to force the Brazilian variety Tupy (an offspring of the Arkansas variety Comanche) to flower and fruit in an area of no chill. The fruit is harvested from mid October until May or early June in this region, and currently provides fresh blackberries for U.S. grocery shelves during the winter months. Mexican production has supplanted Chilean blackberry shipping to the U.S. due to less expensive transportation costs of trucking fruit from Mexico compared to air freight required to move blackberries from Chile. Production area in Mexico is estimated to be 6000-8000 acres. The presence of berries in the marketplace in the winter and late spring enhanced the consumer's awareness of blackberries as a grocery item rather than a local item picked on a farm or from wild plants. The bottom line is that now blackberries are a year-around produce item!

American berry shippers (in the eastern and western US) also took note of the expanded potential of blackberries in the shipping market, spurred by the success

of the Mexican berries shipped and marketed during the “off” season. These marketers felt that if off-season sales could be this successful, why not have expanded marketing during the “normal” US berry production seasons? This has led to an expansion in acreage grown for shipping since the early to mid 2000s, particularly in southern Georgia, Arkansas, Texas, and North Carolina. California greatly increased production in recent years also.

Local production for pick-your-own, farmers markets, or on-farm sales has also increased recently, though it is difficult to determine trends in this area due to few production statistics being available. However, the expanding number of thornless variety options, enhanced fruit quality, and increased interest in berry consumption for human health benefits should positively impact this type of production.

Primocane-Fruiting in Blackberries

The first recorded occurrence of a primocane-fruited blackberry that I am aware of was a wild plant found by L.G. Hillquist of Ashland, Va. There is no record of breeding with this plant until Dr. Jim Moore obtained it in the mid-1960s while accumulating germplasm for the University of Arkansas breeding program. Although primocane fruiting was not pursued for many years in Arkansas breeding, seedlings evaluated in 1997 resulted in Prime-Jim® and Prime-Jan®, released in 2004. Primocane fruiting has been vigorously pursued in Arkansas breeding since the late 1990s, and great headway has been made in improving fruit quality, incorporating thornlessness, and shifting the fruiting period to both earlier and later ripening.

Blackberry Varieties to Consider

Chester Thornless. Although I would like to recommend an Arkansas variety as my top choice for Mid-Atlantic growers, this variety has provided sustained high yields and good hardiness. The main disadvantage of the USDA-ARS-developed Chester Thornless is overall flavor and quality. It ships exceptionally well, but percent soluble solids is not as high as most fresh-fruit consumers desire and a tart taste is normally noted unless fully ripe. This is a semi-erect-caned type. There are other varieties of this cane type such as Hull Thornless, Black Satin, Thornfree, Dirksen Thornless, and Smoothstem, and all are likely adapted to the Mid At-

lantic (they originated in southern Illinois or Maryland), but concerns of tart flavor are often expressed. These varieties tend to be later than Arkansas developments, fruiting in late June to early July in Arkansas.

Triple Crown. The last release of the USDA-ARS varieties, Triple Crown is renowned for exceptional flavor. Some consider this the best-tasting eastern US blackberry. It is moderate to high yielding, appears to have adequate hardiness for the Mid-Atlantic (maybe not quite as hardy as Chester Thornless?), and is earlier in ripening than Chester Thornless (ripens about June 25-30 in Arkansas). The biggest drawback to Triple Crown is berry firmness, and it is not considered a shipping berry. For local markets with short holding times, and pick your own, it is a winner.

Ouachita. If you are considering one Arkansas variety, you should consider this one. It is successful coast to coast in the US, although I have not heard confirmations of its hardiness potential across the entire Mid-Atlantic region. Ouachita produces high yields of high quality berries (6-7 g) with soluble solids of 10-11%. It has erect canes, and ripens about June 10 in Arkansas. It has shown broad adaptation, and has been a major variety in expansion of the domestic shipping blackberry industry.

Navaho. The first Arkansas thornless, Navaho is considered by some to be the best shipping blackberry available. It has medium berry size (5 g) and moderate yield capacity. Sweetness is very good, usually 11-12% soluble solids. Its hardiness has been found to be good in the lower Midwest, and in some areas of the Mid-Atlantic. It is susceptible to orange rust, a fungal disease. It has erect canes, and ripens about June 20 in Arkansas.

Apache. The large-fruited Apache (10-11 g) is admired by some growers, and it has high vigor, productive, and healthy plant characteristics. It averages 11% soluble solids, and ripens about June 25 in Arkansas. Hardiness is not fully known for the Mid-Atlantic, but possibly information exists on this in trials in the region. The major negative attribute of Apache is that white drupes are often seen on some berries, particularly early in the season. This is a very serious defect for shipping, but local sales are usually not impacted as greatly. Concerns among grower reports vary from major to none on this trait.

Natchez. The newest of the Arkansas thornless, Natchez ripens about June 5 in Arkansas. It has large, long berries, and is eye catching on the vine or in the

clamshell. Hardiness is not known on this 2007 release, so care should be taken to determine if it is adapted to the Mid-Atlantic. It averages about 9.5% soluble solids, and berries can be tart if crop load is excessive as it can be in some southern plantings.

Prime-Ark® 45. The first shipping-quality primocane-fruited blackberry released in 2009, it is hoped that Prime-Ark® 45 will provide the basis for developing a late summer to fall-fruited blackberry production season in the US. It has large berries (up to 10 g) with good soluble solids (10% commonly) that stay black in storage along with good firmness retention. The florican crop ripens June 5 in Arkansas, and the primocane crop in mid-August. However, the primocane crop ripe date depends on location. Along the Central Coast of California, first ripe is usually Sept. 1, and in Oregon's Willamette Valley mid-September. This variety has been tested at Penn State Univ. by Kathy Demchak,

and a limitation has been getting good yields before cold temperatures develop. High tunnels have helped, but trials are continuing to determine if adequate yields can be attained in the region. Only trials of the variety are suggested at this time.


Prime-Jan®. Released as one of the first primocane-fruited blackberries in 2004, this variety was originally intended for home garden use. However, limited trials have found it to have some commercial potential. Quality is acceptable, with moderate storage capability and soluble solids on average about 9%. It ripens earlier than Prime-Ark® 45 by about 2 weeks, so has a potential of maturing more of the fall crop prior to frost. However, its crop has not fully ripened (non-high tunnel grown) in upstate New York. Again, testing the variety for specific locations and management (high tunnel or not) is recommended prior to full commercial use is considered.

CALL FOR SPECIALS!

**Reduced prices on
2013 pre-bud orders.
Order early for 2012.**



Shawn Bixby



Debi Clover



Paul Woolley

Quality • Selection • Service

1-800-435-8733



**Stark Bro's
Nurseries &
Orchards Co.**

P.O. Box 398
Louisiana, Missouri 63353
573-754-5111
Fax 573-754-3701
1-800-435-8733

New Jersey News

William E. Johnson Passes at Age 84

Proud, pure, perfect and Republican, WILLIAM EDWARD JOHNSON, passed away on Saturday, September 3rd on his farm in Medford, NJ at the age of 84. Survived by his wife Bette of Medford, his brother Edward of Vermont, sister Maria of Florida, daughter Hille Baldyga and her husband Robert and sons Eric and his wife Karen and Peter and his wife Brenda all of Medford, NJ, Bill was the proud grandfather of nine grandchildren Amanda and Benjamin Baldyga,

Leah, Gilbert, Olivia, Natalie, William S., Wesley, and James Johnson and one great-granddaughter Reese Baldyga. Born in Burlington, NJ, Bill was a life-long resident of Burlington County. On February 14th 1953, Bill and Bette married

after his service in the U.S. Navy and Army. Their love took root on a farm at the corner of Church and Hartford Road in Medford, NJ that Valentine's Day. Once they sold their first ear of sweet corn from a roadside wagon, their business sprouted into the famous Johnson's Corner Farm, a staple for produce and summer picnics in the local area. Many people cannot imagine an autumn without a trip to Johnson's for a wagon ride out to the fields to pick pumpkins

and apples.

Late in life Bill's greatest love became watching people enjoy the farm from his porch as he sat in one of his handmade chairs.

Bill proudly served as a member of a number of civic associations including the NJ State Agricultural Committee, the Burl Co. Board of Agriculture, the Medford-Vincentown Rotary Club, the South

Park Deer Club, Saint Peter's Episcopal Church, the Medford Township Board of Education and the Masons. Bill attained a leadership role in each of these communities. Bill was a devoted member of the unofficial Medport Diner Breakfast Club, which fueled him for a hard



day's work of tending crops and fixing tractors.

Farming is a tough career, but Bill enjoyed every grease stain and bead of sweat that came from his passion for painting the land with produce. Now he's busy paving heaven with golden ears of sweet corn.

In lieu of other expressions of sympathy the family requests memorial donations to the Medford-Vincentown Rotary Foundation, P.O. Box 365, Medford, NJ 08055.

Dave Johnson and His Master Gardener Fruit Team at Rutgers Snyder Farm Make a Difference

Win Cowgill

New Jersey Agricultural Experiment Station, Rutgers Cooperative Extension

At the Rutgers Snyder Research and Extension Farm, we have a large group of dedicated master gardener volunteers from the Rutgers Cooperative Extension programs of Hunterdon, Somerset, and Morris Counties. They have supported our research and extension programs for many years at the farm donating thousands of hours of service.

One group of these extraordinary volunteers is our Master Gardener Fruit Team, lead by Dr. Dave Johnson. Dave and his team have helped in all aspects of the establishment, maintenance, and harvest of my 14 acres of tree fruit research plots at Rutgers Snyder Farm. We would be hard pressed to get all the chores completed without our MG Fruit Team support.

In 2010, one of my NC-140 apple rootstock blocks was discontinued, and we donated the block for use by our Master Gardener Fruit



Team to produce apples for the many food

banks that we support. Dave and his colleagues' pruned and trained this mature block of Gibson Golden Delicious. They hand thinned the fruit, and assisted in removing the perennial weeds. The team replanted the block with new trees to fill in where there was some tree loss. Snyder Farm maintained the block with pesticides, PGRs, and irrigation as needed.



2011 Harvest for our Food Banks Dave's Team Makes a Difference

This week, Dave and his team harvested 5,500 pounds (5 bins) of Gibson Golden Delicious. They were picked into cardboard vegetable bins and were off to the NORWESCAP organization for distribution to the needy the next day. The harvest team consisted of master gardeners Gail Brock, Delores Damiano-Szafran, Charles Rannells, Barbara Harris, Bruce Hoffmann, George Crofts, Kim Goodwin, and Mary

Bowers.

Others on the Fruit Team that have assisted at Snyder are Carl Lewis, John Moskway, Linda Cervino, Linda Lewis, Lisa Mann, Lynn Kocot, and Suzie Delaney.

Rutgers Snyder Farm has donated produce to various food banks and organizations for the past 12 plus years. Last year, Snyder Farm donated 6,000 pounds of apples, and with Dave Johnson's team, we should be able to double that to 12,000 lbs in 2011!

Ed Dager, Snyder Farm Supervisor, coordinates all of the donations to the various food banks.

Food Banks

The following are food banks and organizations to which Rutgers Snyder Farm donates annually. These organizations are always looking for additional donations and support to feed the hungry and help those in need.

NORWESCAP (<http://www.norwescap.org>). The NORWESCAP Food Bank aids in the relief of hunger by providing food and support to non-profit charitable organizations in Hunterdon, Sussex, and

Warren Counties. They distribute over 2,000,000 pounds of food annually to over 120 pantries, shelters, soup kitchens, on-site feeding programs, childcare centers, senior centers, and programs for the disabled.

Giving Gardens at Wagner Arboretum Foundation (<http://www.wfafnj.org/ggp.html>). Volunteers supply food and produce to the needy in Somerset County through appropriate agencies.

Mercer Street Friends (<http://www.mercer-streetfriends.org/>). Opened in 1958 by Quakers and promulgating the values of tolerance, equality of all people, social justice, and nonviolent resolution of conflict, Mercer Street Friends works tirelessly to touch the lives of more than 30,000 people a year and help them overcome the debilitating effects of poverty, hunger, neglect, and health concerns. This is about two-thirds of the people in poverty in Mercer County.

The Quakertown United Methodist Church, (Route 579, Quakertown, New Jersey) and ***Frenchtown Presbyterian Church*** (22 4th Street Frenchtown, New Jersey, <http://fpc.frenchtownpres.org/>) distribute produce to needy individuals weekly.





Horticultural News

New Jersey State Horticultural Society
P.O. Box 116
Clayton, NJ 08312