

Horticultural News



Volume 93, Number 1
Winter, 2013

Horticultural News

Editors: Winfred P. Cowgill, Jr. & Wesley R. Autio

The New Jersey State Horticultural Society was organized on August 17, 1875 at Geological Hall, Rutgers College, New Brunswick, NJ. It remains the oldest Horticultural organization in New Jersey.



Horticultural News began as the *The New Jersey State Horticultural Society News*, in October of 1920. The Society began “collecting paid membership in order to obtain funds to promote new features of the society and extend the usefulness of the society. The Horticultural Society News was started to be the official society publication.” Published M. A. Blake, Professor at Rutgers College was the first president and chair of the publication committee.

Editors served as follows:

MA Blake	1920 - 1947
Norman F Childers	1948 - 1980
Win Cowgill	1981 - 1988
Emily Brown Rosen	1988 - 1990
Linda Butenis Vorsa	1991 - 1995
Jerry Frecon	1995 - 2010

June 2010: *Horticultural News* has moved to an online web-based format. The New Jersey State Horticultural Society has partnered with the University of Massachusetts *Fruit Notes*, Dr. Wesley Autio, Editor. Cowgill and Autio will be the new editors of *Horticultural News* and *Fruit Notes*.

Horticultural News is distributed to growers, extension personnel and researchers and libraries across North America. **Horticultural News** focuses primarily on tree-fruit culture, but addresses small-fruit cultural issues as well. Most reports are from current research at Rutgers University, University of Massachusetts, and other universities.

Horticultural News is published four times per year by the New Jersey State Horticultural Society. It is provided as a benefit to membership in the society. Membership costs \$50 per year. Each one-year subscription begins January 1 and ends December 31. Payments via check must be in United States currency and should be payable to the New Jersey State Horticultural Society. **Horticultural News** Electronic subscriptions are available as benefit of membership in the NJ State Horticultural Society. A hidden link will be mailed immediately after publication to members. Issues will be made freely available on this website six months after publication.

Correspondence should be sent to: New Jersey State Horticultural Society
P.O. Box 116
Clayton, NJ 08312
TEL (856) 207-7123

For advertising opportunities, contact: Win Cowgill: wincowgill@mac.com; (908) 489-0207

Table of Contents

Blossom Thinning Results in an Early Bloom Season Tara Auxt Baugher, James Schupp, Catherine Lara, and Sladjana Prozo.....	1
Pesticide Reduction with Containment Spraying of High-density, Dwarf Apple Orchards Wesley Autio, James Krupa, and Jon Clements	10
Appletesters.net -- A new website to assist apple growers and apple testers in variety selection Win Cowgill and Jon Clements	16
Dr. Anne Nielsen, New Fruit Entomology Joins Rutgers Cooperative Extension	20
2013 Mid-Atlantic Convention Program, January 29-31, Hershey, PA.....	22
New Jersey News.....	25

Cover: Darwin 300 String Thinner (designed Fruit-Tech) for vertical apple canopies. Photo by Loy Elliott.

Blossom Thinning Results in an Early Bloom Season

Tara Auxt Baugher, James R. Schupp, Catherine Lara, and Sladjana Prozo
The Pennsylvania State University

Hand thinning is a necessary and costly management practice in peach production. Stone fruit producers are finding it increasingly difficult to find a workforce to manually thin fruit crops, and the cost of farm labor is increasing. The conventional method for adjusting crop load in peach and nectarine orchards is to remove excess fruit by hand at 35 to 40 days after full bloom. Plant growth regulators are available for thinning pome fruit; however, chemical thinning options for stone fruit are limited and unpredictable.

In mechanical blossom thinning trials conducted over five seasons, string thinner crop load management technologies (Figure 1) were tested in four peach producing states, and detailed research on pruning modifications and application timing provided information to guide producers in maximizing mechanical bloom thinning benefits. The original string thinner evaluated in 2007 (Darwin 300, Fruit-Tec, Deggenhausertal, Germany; Schupp et al., 2008) was designed to thin narrow vertical apple canopies and therefore was evaluated on peach trees trained to either a perpendicular V or quadrilateral V system. A prototype designed in 2008 was successfully tested to operate in a horizontal position for thinning trees trained to an open-center system (Baugher et al., 2009). A “hybrid” string thinner (PT250; Figure 1) designed to adjust crop load in either vase or angled tree canopies was evaluated in fresh fruit and pro-

cessing plantings in 2009 to 2011.

String thinner trials with variable tree forms uti-



Figure 1. Blossom thinning in peach with a prototype Fruit-Tec PT250. Photo by Mark Wherley.

lized by producers in California, Washington, South Carolina, and Pennsylvania demonstrated reduced labor costs compared to hand-thinned controls and increased crop value due to a larger distribution of fruit in marketable and higher market value sizes (Baugher et al., 2010a). Blossom removal ranged from 17% to 56%, hand thinning requirement was reduced by 19% to 100%, and fruit yield and size distribution improved in at least one string thinning treatment per experiment.

Research in Pennsylvania orchards was conducted over two years to evaluate string blossom thinner efficacy at variable stages of bloom development, ranging from pink to petal fall (Baugher et al., 2010b). Blossom removal at the pink stage of bloom development was lower than at other stages in 2008; however, a 150 rpm versus 120 rpm spindle rotation speed resulted in blossom removal similar to a 80% full bloom treatment in 2009. Blossom removal at the petal fall stage was similar to the open bloom stage. Savings in hand thinning time and/or increases in fruit size in both years associated with the bloom stage treatments resulted in a net positive impact of \$49 to \$554 per acre compared to hand thinning alone.

Pennsylvania studies also were conducted over two seasons in peach orchards trained to perpendicular V or open center systems to evaluate possible pruning strategies to improve tree canopy access by string thinners (Schupp et al., 2011). The objectives were to demonstrate if modifications in fruiting shoot orientation, pruning detail, and/or scaffold accessibility improved flower removal, reduced follow-up hand thinning requirement, and/or increased fruit size. Blossom removal was improved by either detailed pruning (elimination of short or excessively long shoots) or partial pruning (elimination of all shoots on the side of a limb inaccessible by the thinner spindle) in both training systems. The best treatments resulted in a thinning savings of \$49 to \$282 per acre in perpendicular V plantings and \$11 to \$19 per acre in open center plantings.

Case study interviews of 11 Pennsylvania growers and orchard managers who had thinned a total of 154 acres suggested that commercial adoption of mechanical string thinning technology would have positive impacts on the work place. All case study cooperators reported that blossom string thinning impacted orchard management by making crop load management more efficient and by reducing follow-up hand thinning time. Eighty percent of the growers noted

fruit from thinned trees were larger. Additional observations included the following: 1) hand thinning of peaches was completed earlier allowing more timely work in other crops, 2) employees were satisfied with mechanical thinning as it saved them time and minimized ladder use, and 3) the seasonal distribution of labor-intensive work was improved.

What about Thinning during an Early Bloom Season?

One lingering question that producers considering bloom thinning have had is “Should we bloom thin in an early season when the potential for freeze injury may be greater?” The hypotheses tested were: 1) bloom thinning in an early season should only be conducted on cultivars that will withstand some additional thinning from freeze injury, and 2) string thinner spindle rotation speed should be reduced in a year when there may be more potential for freeze injury.

Materials & Methods

In 2012, South-central Pennsylvania peach orchards began to bloom four weeks ahead of the normal timing (early March vs. mid-April). Many producers kept their string thinners in their equipment sheds, but two growers agreed to participate in trials in open-center trained orchards with two early maturing cultivars for which optimizing fruit size is important—‘Rising Star’ and ‘Glenglo’—and in two ‘John Boy’ blocks that tend to be reliable producers. In each of the four trials, two string thinner spindle rotation speed treatments were compared to control treatments that were hand thinned at the green fruit stage. Flowers/fruit were counted before thinning, during the physiological drop stage, and prior to follow-up hand thinning. Follow-up hand thinning at the green fruit thinning stage was conducted on the rpm treatments to assess effects on labor requirement. Fruit were measured at harvest to assess effects on fruit size. The plots were arranged in randomized complete block designs with six multiple tree replicates. Data were collected from center trees and subjected to analysis of variance. Labor costs were provided by cooperating growers, and machine costs were obtained from equipment manufacturers. Peach market values for various size categories were obtained from the USDA Agricultural Marketing Service report for the Appalachian region (USDA, 2012).

Results & Discussion

Peach Blossom Thinning and Fruit Set Response in a Year with Increased Potential for Freezing Temperatures during Bloom. Initial flower density ranged from 15.3 to 25.1 flowers per cm² limb cross-sectional area across the four orchard plots. Flower density was reduced by thinning treatments in two of the four peach orchard plots (Table 1). In the plots where flower density was not reduced (Orchard A), flower removal ranged from 6.8% to 22%; whereas in the plots where flower density was reduced, the grower (Orchard B) had selected rpm treatments that removed 42% to 61% of the blossoms. In ‘John Boy’, Orchard B, the 220 rpm treatment removed more flowers than the 200 rpm treatment; however flower removal in ‘Rising Star’ was equal in 175 and 200 rpm treatments. Prior to the green fruit thinning stage, crop load was more than desired across all treatments and all required follow-up hand thinning.

There were two freeze events prior to thinning (low temperatures of 29° to 32°F) and three freeze events following bloom thinning (low temperatures around 32°F) (Figure 2, minimum temperatures from weather station at Penn State Fruit Research and Extension Center, Biglerville, PA). The freeze events prior to thinning reduced crop load by approximately 10%. In Orchard B, percent change in flowers/fruit remaining from the dates of bloom thinning to fruit set was significantly higher in the control treatments than in the string thinning treatments. The reduced natural drop might be explained by the reduced competition between fruitlets, which may provide a “cushion” in years with an increased possibility of freeze injury. During the prior six years during which we conducted bloom thinning studies in Pennsylvania orchards, there was one trial that was subjected to freezing temperatures following bloom, and a similar trend was observed. In this case, flower density was reduced by 90% by freezing temperatures, but at fruit set, the crop load in string thinned treatments was equal to that in the control treatment (Baughner et al., 2010b). Since temperatures did not drop below 32°F in the current study, we cannot draw conclusions about critical temperature events that kill 90% of blossoms.

Mechanical Blossom Thinning Effects on Labor Requirement and Fruit Size. Hand thinning in Orchard B’s ‘Rising Star’ and ‘John Boy’ was reduced by all bloom thinning treatments, and the thinning savings ranged from \$25 to \$48 per acre. Although

the crop load comparisons in Orchard A were non-significant, hand thinning of ‘John Boy’ was reduced by the 210 rpm treatment, which resulted in a \$36 savings per acre. The loss in the other Orchard A treatments was \$15 per acre—the cost of mechanical thinning.

Fruit diameter was improved in the 200 rpm treatment in ‘Rising Star’ and the 200 and 220 rpm treatments in Orchard B ‘John Boy’. Fruit in the higher value 2 ¾ inch and higher fruit size categories was increased in both ‘John Boy’ plots and in ‘Rising Star’. As the season progressed, fruit of variable sizes and shapes were observed in the two early season cultivars, which were probable effects of sub-lethal temperature injury and pollination conditions interacting with genetics, and harvestable yields were reduced across the bloom thinning treatments and the controls. In these plots, the net impact per acre of bloom thinned compared to hand-thinned control treatments (taking into account effects on labor requirement and fruit size) ranged from -\$15 per acre for ‘Glenglo’ to \$171 per acre for the 200 rpm treatment in ‘Rising Star’. By comparison, value added by increases in fruit size, increased in both ‘John Boy’ plots, and net impact ranged from \$619 to \$1624 per acre, which is consistent with the impacts in prior research conducted in more normal bloom seasons (Baughner et al., 2010a, 2010b; Schupp et al., 2009, 2011).

Conclusions. In a growing season that began four weeks early, temperatures dropped to freezing levels on three occasions but did not reach critical lows. Therefore, the hypotheses could not be fully tested. Across four research plots, the thinning effects on a reliable producing cultivar were generally positive, but effects on early maturing cultivars were variable. The higher compared to the lower rpm resulted in more fruit in higher value size categories in two of the four trials but equal reductions in follow-up hand thinning requirement. The economic impacts from the increases in fruit size were \$146 and \$562 per acre greater for the higher rpm treatments in ‘Rising Star’ and Orchard B ‘John Boy’, respectively. The question of whether or not to bloom thin in an early bloom season will remain a question to be addressed for specific orchard blocks based on site history and cultivar susceptibility, but the early 2012 season provided an opportunity to obtain some guiding information. As producers gain experience with optimum spindle speed in various cultivars and sites in a normal bloom season, they will learn how to adjust rpm for a year in which crop potential may be reduced.

Table 1. Peach blossom thinning and fruit set response to mechanical thinner treatments applied in an early growing season.

Cultivar/Orchard	Treatment ^z	Flower removal (%)	Flower density before/after thinning (flowers/cm ² LCSA ^y)	Crop load (fruit set) 30 DAFB ^x (fruit/cm ² LCSA)
Glenglo Orchard A	Hand-thinned control			
	30 DAFB	--	25.1/25.1 a	8.3 ab
	String thinned 150 rpm	6.8 b ^w	20.5/17.1 a	11.3 a
	String thinned 180 rpm	18.4 a	21.5/15.6 a	6.4 b
John Boy Orchard A	Hand-thinned control			
	30 DAFB	--	19.3/19.3 a	9.8 a
	String thinned 180 rpm	15.2 a	17.1/14.4 a	7.9 a
	String thinned 210 rpm	21.6 a	20.1/15.7 a	7.6 a
Rising Star Orchard B	Hand-thinned control			
	30 DAFB	--	11.8/11.8 a	9.3 a
	String thinned 175 rpm	50.7 a	12.2/6.6 b	5.5 ab
	String thinned 200 rpm	61.1 a	10.6/4.4 b	3.6 b
John Boy Orchard B	Hand-thinned control			
	30 DAFB	--	17.8/17.8 a	11.2 a
	String thinned 200 rpm	42.2 b	15.3/8.4 b	6.5 b
	String thinned 220 rpm	61.0 a	17.5/6.9 b	5.0 b

^z Peach trees were thinned at 50 to 100% full bloom. Tractor speed was 2 mph.

^y Limb cross-sectional area.

^x Crop load assessed just prior to hand thinning at the green fruit stage. DAFB = Days after full bloom

^w Mean separation within columns and cultivars by Fisher's protected least significant difference at $P \leq 0.05$.

Literature Cited

Baughner, T. Auxt, J. Schupp, K. Lesser, and K. Reichard. 2009. Horizontal string blossom thinner reduces labor input and increases fruit size in peach trees trained to open-center systems. *HortTechnology* 19:755-761.

Baughner, T. A., J. Schupp, K. Ellis, J. Remcheck, E. Winzeler, R. Duncan, S. Johnson, K. Lewis, G. Reighard, G. Henderson, M. Norton, A. Dhaddey, and P. Heinemann. 2010a. String blossom thinner designed for variable tree forms increases crop load management efficiency in trials in four United States peach-growing regions. *HortTechnology* 20:409-414.

Table 2. Follow-up hand thinning required for mechanical thinner treatments applied in an early growing season.

Cultivar/Orchard	Treatment ^z	Hand thinning at 30 to 35 DAFB (h/acre/1 person)	Thinning savings ^z (\$/acre)
Glenglo Orchard A	Hand-thinned control 30 DAFB	28.2 a ^y	--
	String thinned 150 rpm	28.8 a	(15)
	String thinned 180 rpm	29.7 a	(15)
John Boy Orchard A	Hand-thinned control 30 DAFB	26.8 a	--
	String thinned 180 rpm	24.5 ab	(16)
	String thinned 210 rpm	22.8 b	36
Rising Star Orchard B	Hand-thinned control 30 DAFB	9.9 a	--
	String thinned 175 rpm	7.2 b	25
	String thinned 200 rpm	6.1 b	35
John Boy Orchard B	Hand-thinned control 30 DAFB	17.9 a	--
	String thinned 200 rpm	12.5 b	48
	String thinned 220 rpm	13.4 b	41

^z Thinning savings includes reduced follow-up hand thinning inputs and added mechanical thinner, tractor, and labor inputs. Values in parentheses are negative and represent cost of mechanical thinning.

^y Mean separation within columns and cultivars by Fisher's protected least significant difference at $P \leq 0.05$.

Table 3. Peach fruit size, high value packout distribution, market value based on fruit size, and net economic impact from both labor savings and/or fruit size increase.



Cultivar/Orchard	Treatment	Fruit diameter ^z (cm)	Fruit $\geq 2 \frac{3}{4}$ inch ^z (%)	Added value (\$/acre)	Net impact (\$/acre) ^y
Glenglo Orchard A	Hand-thinned control				
	30 DAFB String thinned 150 rpm	7.4 a	69 a	--	--
	String thinned 180 rpm	7.4 a	60 a	--	(15)
John Boy Orchard A	Hand-thinned control				
	30 DAFB String thinned 180 rpm	7.4 a	48 b	--	--
	String thinned 210 rpm	7.4 a	60 a	635	619
Rising Star Orchard B	Hand-thinned control				
	30 DAFB String thinned 175 rpm	7.7 b	79 b	--	--
	String thinned 200 rpm	7.7 b	79 b	---	25
John Boy Orchard B	Hand-thinned control				
	30 DAFB String thinned 200 rpm	7.5 b	40 b	--	--
	String thinned 220 rpm	7.7 a	55 ab	1015	1062
		7.8 a	79 a	1584	1624

^z Fruit diameter and high value packout distribution determined on 40 fruit harvested per treatment from each of six replicates.

^y Net economic impact (realized economic savings) is defined as cost or benefit beyond hand thinning alone and takes into account reduced hand thinning inputs and increased value of fruit in higher size categories. Values in parentheses are negative.

^x Mean separation within columns and cultivars by Fisher's protected least significant difference at $P \leq 0.05$.

Figure 2. Crop load and low temperature mean comparisons from bloom to fruit set (minimum temperatures from weather station at Penn State Fruit Research and Extension Center, Biglerville, PA).



Baughner, T. Auxt, J. Schupp, K. Ellis, J. Remcheck, E. Winzeler, K. Lesser, and K. Reichard. 2010b. Mechanical string thinner reduces crop load at variable stages of bloom development of peach and nectarine trees. *HortScience* 45:1327-1331.

Schupp, J.R., T. Auxt Baughner, S.S. Miller, R.M. Harsh, and K.M. Lesser. 2008. Mechanical thinning of peach and apple trees reduces labor input and increases fruit size. *HortTechnology* 18:660-670.

Schupp, J. R. and T. Auxt Baughner. 2011. Peach blossom string thinner performance improved with selective pruning. *HortScience*

46(11):1486-1492.

U.S. Department of Agriculture. 2012. USDA agricultural marketing service report. USDA fruit and vegetable market news. Aug. 2012 <http://marketnews.usda.gov/portal/fv?paf_dm=full&paf_gear_id=1200002&startIndex=1&dr=1&rowDisplayMax=25&repType=termPriceDaily&dr=1&locName=&comAbr=PCH&commName=PEACHES>

Acknowledgements

The authors acknowledge the valuable contributions of Dave Wenk, Eddie Rankin, and Arturo Dias.

PEACHES

Still growing strong!


**Delaware & California Grown
Certified Peach Trees.
Order Now for Spring.**

ACN INC. SINCE 1905

**Adams County Nursery, Inc.
Aspers, PA
(800) 377-3106 • (717) 677-4124 Fax
Website: www.acnursery.com
Email: acn@acnursery.com**

Be on the Leading Edge ...
RubyMac®
USPP # 19,891 • An IPM Variety

- Blushed, vibrant ruby-red finish over virtually 100% of the fruit.
- Colors weeks ahead of other Macs
- Outstanding firmness and quality
- Ripens in Mac season with great flavor
- Consistently delivers a high pack-out



An IPM Variety.

“This is a phenomenal Mac. Nothing out there compares. It blows the doors off every other Mac in the field. It colors early and finishes beautifully. We expect a much higher pack-out with RubyMac™. The real beauty of RubyMac™ is that we don't have to wait for color and sacrifice condition. We can pick it without waiting for color and we have a firmer, better Mac that delivers a higher quality product to our customers. **It's the Mac we've always wished we had!**”

— Pat Chase
buyer for Jack Brown Produce, Inc.
Sparta, MI



SUMMIT TREE SALES
REPRESENTING THE COUNTRY'S FINEST NURSERIES

800.424.2765 / 269.674.8866 / Fax 269.674.3758
summit@summittreesales.com

Pesticide Reduction with Containment Spraying of High-density, Dwarf Apple Orchards

Wesley R. Autio, James S. Krupa, and Jon M. Clements
University of Massachusetts

Project Purpose

With the adoption in commercial apple orchards of fully dwarfed trees and ultra-high-density planting/training systems, containment or tunnel spraying becomes a feasible alternative to conventional airblast spraying. By spraying only within a moving canopy extending from the sprayer on both sides of the row being treated, tunnel spraying can dramatically reduce drift of agricultural chemicals to non-target areas and substantially decrease the quantity of chemical required. The potential environmental benefits are unmeasured. The quality of pest control should be enhanced, while at the same time, significantly reducing the quantity of pesticide used. A change in technology such as this can only enhance the sustainability and competitiveness of apple farming in a steadily urbanizing part of the US.

Through this project, the Massachusetts Fruit

Growers' Association (MFGA) and the University of Massachusetts Fruit Program (UMass) addressed the following objectives: (1) demonstrate the feasibility of tunnel-sprayer technology in Massachusetts orchards; (2) estimate drift to demonstrate whether or not this approach will reduce environmental risk within and near orchards; (3) assess efficacy of reduced chemical application rates per acre in an effort to adjust rates and recommendations to account for much smaller tree volume per acre.

Summary of Results

The Lipco Tunnel Sprayer was delivered to the UMass Cold Spring Orchard on October 1, 2010. Hans Wörthle from H&W Equipment visited on October 19 and 20, along with a crane, to assemble the tunnel sprayer. It was tested briefly and then winterized.

Significant work with the sprayer began in April 2011. Because of the dramatic differences between it and conventional sprayers, it took time to become familiar with its operation and manipulation. First observations were: 1) the sprayer is very tall and because it is offset behind the tractor, it requires more care when driving down a tree row; 2) a wind parallel to the row can blow drift out of the front or back of the sprayer; and 3) the tunnel sprayer is much quieter than an airblast sprayer.

Using published charts relative to the fluid flow out of the sprayer nozzles, we adjusted the tunnel sprayer and an airblast



Figure 1. Lipco Tunnel Sprayer at the UMass Cold Spring Orchard Research & Education Center.



Figure 2. Rear view of a typical airblast orchard sprayer.

be approximately 10-20% of the spray material; this amount would be much larger on a windy day. The tunnel sprayer, however, produced no measurable drift.

To measure the relative effectiveness of spraying with a tunnel sprayer versus a conventional airblast sprayer, a study was conducted in 2012, comparing the applications of two nutrient sprays with each sprayer. A block of approximately 200 Silken trees that were trained to a tall-spindle system was used for this trial. Trees were divided among six replications of an experiment including an untreated control and calcium chloride (at the recommended rate) and an experimental formulation of calcium from Key-Plex



Figure 3. Even under nearly calm conditions, the airblast sprayer has the potential to create spray drift if it is not well calibrated.

applied with the tunnel sprayer or with the conventional airblast sprayer. Treatments were applied three times throughout the summer. Leaf samples and fruit samples were collected at the end of August. Leaf samples were submitted to the UMass Soil & Tissue Analysis Laboratory for the assessment of nutrient element concentrations. The fruit samples were submitted to the Fruit Program's Fruit Analysis Laboratory for the assessment of calcium concentrations. Unfortunately, fruit tissue results were not yet available at this writing. Leaf analyses, however, showed no significant differences between the types of sprayers, and

sprayer to deliver the same amount of material per acre. Observation suggested that the airblast sprayer resulted in much more drift, but coverage appeared better than with the tunnel sprayer. This observation puzzled us, so we measured flow out of all of the nozzles and found that the published flow rates were wrong. To obtain the desired flow, we purchased new nozzles, and selected air-induction nozzles (to increase particle size and reduce drift potential). With the new nozzles, the tunnel sprayer provided excellent coverage, with far less drift than the airblast sprayer.

Drift (utilizing water-sensitive paper) was measured on a reasonably calm day. The airblast sprayer, although calibrated well, produced some drift beyond the target trees. It was estimated to



Figure 4. Airblast orchard sprayer in operation. Spray material clearly passes through the canopy. Likely, most will fall on the trees in the next row, but if weather conditions are undesirable, this material may move into non-target areas.



Figure 5. The Tunnel Sprayer directs material into the canopy from the outside and captures that which passes through the canopy.

Hands-on demonstrations were conducted at twilight meetings on May 17, 2011, and April 17, 2012, at the UMass Cold Spring Orchard Research & Education Center (35 and 30 farmers in attendance in 2011 and 2012, respectively), and presentations (with video) were given at three additional twilight meetings (May 18, 19, and 26, 2011) with total attendance of 129 farmers. It also was demonstrated at the 2012 Annual Summer Meeting of the Massachusetts Fruit Growers' Association at the UMass Cold Spring Orchard Research & Education Center on July 16, 2012, with approximately 100 farmers in attendance. Small-scale demonstrations were conducted several times during the two years to a total of approximately 200 individuals.

In total, about 350 people have seen a hands-on demonstration of the tunnel sprayer. About 130 have seen presentations given in person with video demonstrations, and another 4,700 have viewed web-based presentations.

the expectation is that there likewise will be no differences in the fruit samples.

Educational programs began in earnest in 2011. Details of this project have been recorded in a blog: <http://masscon.blogspot.com> (Massachusetts Containment Spraying Blog). Four video presentations are provided in the blog to describe progress during the early stages of the process. The blog has been visited a total of 1,579 times since its creation 16 months ago. The videos were also provided on YouTube (<http://www.youtube.com/user/wrautio1>) and, in total, have been viewed 3,142 times.



Figure 6. Water sensitive paper showing drift from the airblast sprayer (left photo) and the lack of any spray drift from the tunnel sprayer (right photo).




Figure 7. The drift trial measured drift onto water sensitive paper at 1, 5.5, and 10 feet from the soil surface on posts 10, 20, 30, and 40 feet from the outer tree row.

Outcomes and Lessons Learned

1. *Sprayer is feasible under our orchard conditions. Hydraulic manipulation of the height and widths of the sprayer allows adjustment specific to tree size and slope.*
2. *Spray use is reduced by 10 to 20%, likely with no change in efficacy (the latter point still needing full verification).*
3. *Drift is nearly nonexistent with the tunnel sprayer; whereas, it represents 10-20% of the spray material with an airblast sprayer. Environmental benefits are not easily measured but could be significant. Likewise, eliminating drift may allow farming closer to humans, without any risk of off-target exposure.*
4. *This technological advancement comes at a cost of about \$30,000 per sprayer. The extra cost cannot be offset by the cost benefit of reduction in spray material. The potential environmental benefits must be weighed before deciding to purchase such a sprayer. In some settings, it may be becoming impossible to use airblast technology because of the proximity to human dwellings, and this sprayer provides an alternative.*
5. *Overall, the sprayer worked very well, performing exactly as expected. We cannot recommend it to the general grower because of price; however, we can recommend it under situations where drift is an insurmountable problem.*

Acknowledgements

This project was funded by a USDA Specialty Crops Block Grant through the Massachusetts Department of Agricultural Resources. The authors are very thankful for this support.

RIGHT...FROM THE START

Start your orchard RIGHT with Willow Drive Nursery's Proven Performers™. Call for a complete and updated inventory.

Calling us today can help you tomorrow!



APPLES

Braeburn

Mariri Red™ (Eve®)
Kumeu Crimson®

Fuji

Aztec Fuji® (DT 2 Variety)
Rising Sun®
Morning Mist™
Early Fuji

Gala

Brookfield® Gala
Buckeye® Gala
Ultima Gala®

Other

Cameo® brand
Granny Smith
Honeycrisp™

It® Red Delicious
Morren's® Jonagored
Supra™
Ruby Jon®
RubyMac®
Smoothee® Golden
Spartan



CHERRIES

Attika®

Benton™

Bing

BlackGold™

BurgundyPearl™

Chelan™

Early Robin®

EbonyPearl™

Lapins

Montmorency

Rainier

Regina

Rynbrandt

Schmidt

Selah™

Skeena™

Sweetheart™

Tieton™

Ulster

WhiteGold™



PEARS

Bartlett

Columbia Red Anjou™

Concorde™

Comice

D'Anjou

Forelle

Golden Russet Bosc®

Red Clapp's Favorite



PEACHES

Stellar Series

Allstar®

Autumnstar®

Brightstar™

Coralstar®

Earlstar™

Flamin' Fury

PF 8 Ball

PF Lucky 13

PF 19-007

PF 25

PF 35-007



WILLOW DRIVE NURSERY, INC.

1-888-54-TREES

Ephrata, Washington · 1-888-548-7337 · Web Site: www.willowdrive.com

Targit Sales

“Specializing in Farm Market Packaging”
1-800-526-9224

Biodegradable

PLASTIC BAGS



Targit Sales is proud to introduce our line of Biodegradable plastic bags. Our bags use a proprietary additive that renders the plastic ultimately biodegradable.



We are offering the Biodegradable bag program with all of our “Custom Imprinted” plastic bags. Our Biodegradable bags have the “EPI” (the additive company) logo on the bag as proof of its biodegradable properties.

Advertise your business along with a “tag line” to express your company’s efforts to keep your community and planet green.

Targit Sales can fully customize your bags to your specifications. We can make any size, strength, and style of bag to fit your needs. We can assist you with your design, setup, and artwork.

COMMON SIZES OF BAGS:

- 13 x 10 x 24 (Great for pies, cakes and corn)
- 12 x 8 x 24 (Great for your store front)
- 8 x 5 x 21 (Perfect size for tailgate markets)

BIODEGRADABLE PLASTIC BAG BENEFITS!

- * Superior Strength!
- * Less expensive to make and ship versus paper bags!
- * Superior imprinting capability and quality...(looks better)!
- * Easier and more affordable to store (inventory)!
- * Flexibility! (customize the size, thickness, color and style to your specifications)
- * Reusability!
- * Biodegradable..breaks down quickly!

MOST COMMON QUESTIONS asked about biodegradable bags.

Will the bags begin to break down while being stored? No, the bags will retain its strength and integrity while properly stored. The breakdown only begins after it is directly exposed to certain environmental conditions such as: soil/landfill, heat/moisture, and sunlight.

Is this process more expensive? Not really, the cost increase is only a few percentage points higher. You will more than make this up with the goodwill you will be conveying to your community and customers.

Does the bag lose its strength or feel...does it look different? No, it is almost impossible to tell the difference between the Biodegradable bags and the traditional plastic bags. This is why we recommend putting a tag line on your printed biodegradable bags.

Can these bags be recycled and/or reused? Yes, these bags can be recycled and reused.

For more information or to order contact
Targit Sales
1-800-526-9224 • www.targitsales.com



Appletesters.net – A New Website to Assist Apple Growers and Apple Testers in Variety Selection

Win Cowgill

New Jersey Agricultural Experiment Station

Jon Clements

University of Massachusetts Extension

Appletesters.net is a new website designed to assist apple growers in variety and cultivar selection for new orchard plantings. It is also designed to provide a location for anyone conducting cooperative apple evaluation and testing to report and archive their findings using a unified protocol for apple variety fruit and trees.

Variety and cultivar selection is the largest economic decision a grower can make when establishing a new high-density orchard block of a thousand trees or more per acre using tall spindle systems.

Appletesters.net allows growers or potential growers to log on and view records of a variety from different locations over multiple years to assist making their planting decisions.

The online records allow cooperators and growers to see what variety characteristics are performing best in their area. The database will include pictures and casual observations in addition to fruit quality measurements and important horticultural characteristics including tree habit, bloom time, disease tolerance, mortality, precociousness, fruit appearance, taste, storability, and more.

Directions for Use

Point your web browser to: <http://appletesters.net>. Click on the apple testers database Guest Login. This brings you to our login page, select the guest account button and click login. This brings you to a data base record; you are now in the system. The first thing you should do is familiarize yourself with the database layout. In the tool bar at the top, all the way

to the right is a help button. Click the help button to open a new window that contains the directions as to how the database works. Read through and see what it's all about. This window will stay open for reference for you at any time.

Example of Use

Lets look at one search as an example of one way to use the database. In the tool bar select the button Find. This will bring up a blank data base record, you can search on any criteria but lets select the cultivar box by typing a cultivar/variety, lets say Zestar. Then go back up to the tool bar and click the Perform Find button. This brings up 7 records for Zestar. You know this by again looking at the toolbar on top in the left hand corner at Found Records. With Zestar (As of 12/15/12) you will see 7 records that you can cycle trough using the arrows to the left of the Found Records icon.

Appletesters.net will become an important repository of cultivar/strain information on tree growth and fruit quality over many seasons. Such information should be very useful to apple growers looking for more and/or unbiased variety/strain information to make future planting decisions.

Funding

Funding for this project was provided by a North East SARE partnership grant.
<http://www.nesare.org/>.



State MA
Location Belchertown
Cultivar Paulared
date 8/19/2011
fruit picture <http://fruit.umext.umass.edu/2011apple/paulared0819.jpg>



Figure 1. Apple Tester database picture of Paulared record, Belchertown, MA 8/11.

appletesters

http://128.119.96.130/fmi/iwp/cgi?-db=applete: Google

APS - Journal Growing Wit...opean Flair hazecam Facebook UMass Fruit Advisor

appletesters AppleTesters 2010 report

4 72 / 317 Found (Unsorted) Records Show All New Record Edit Record Delete Record Records Find

Layout: appletesters View As: [Table Icon] [List Icon] [Form Icon]

appletesters.net

date 8/19/2011

State MA

Location Belchertown rootstock

Cultivar Paulared

tree vigor medium-strong

branching medium

TREE CHARACTERISTICS tree habit spreading-drooping

crop load medium-heavy biennial

maturity period early

pre-harvest drop some

fruit ground color green

fruit over color red

color % red 75

color pattern blush

lenticels inconspicuous

FRUIT APPEARANCE fruit symmetry globose

russet coverage nil

russet region

russet texture

cracking absent

cracking location

fruit size medium-large

fruit diameter inches 3.1

fruit weight grams 184

firmness lbs 15.1

soluble solids 11

starch index 3.5 (2-5) browning faint

flesh color green-white

flesh texture soft

flavor tart

FRUIT FLAVOR skin thickness average

skin greasiness dry

eating quality fair

Figure 2. Blank Apple Tester Database Record Form.

LET
OESCO
POWER
YOU
THROUGH
THE
GROWING
SEASON!

For more information
visit our
NEW website at
www.oescinc.com



REARS POWERBLAST
A powerful machine with PTO drive and the patented Constant Velocity Hitch (CVH). 400 or 500 gallon tank.



REARS PAK TANKS
3 Pt. hitch, pto drive diaphragm pumps, choice of booms or guns.



REARS PUL BLAST
100-500 gallon tanks. Medium sized, variable pitch fans. Diaphragm or centrifugal pumps.



**REARS FLAIL MOWER
OMF-750 SERIES**
Super-duty low profile, uses a double row spherical roller bearing with oversize rollers and eccentric locking collar.



8 Ashfield Road on
Route 116
Conway, MA 01341

800-634-5557 • 413-369-4335 • info@oescinc.com

**Need Used Equipment—
We Have That Too!
CALL US FOR DETAILS.**



80
YEARS

Since 1932

The Best Berry Plants

- Strawberries, raspberries, blueberries, blackberries, asparagus and more!
- Where the pros go for plans and plants.
- Call for a free catalog and plasticulture guide!

41 River Road, South Deerfield, Massachusetts 01373


NOURSE

www.noursefarms.com 413.665.2658

Dr. Anne Nielsen, New Fruit Entomologist Joins Rutgers Cooperative Extension

Anne L. Nielsen is the fruit entomology extension specialist located at RAREC in Bridgeton, NJ. Her responsibilities include tree fruit, small fruit (excluding blueberries and cranberries) and grapes. Dr. Nielsen joins Rutgers after postdoctoral research experiences at UC Davis and Michigan State University. Happy to be returning to Rutgers, where she completed her dissertation under the direction of George Hamilton, Anne's research will focus on developing and refining phenological models for key pest species. This basic research is combined with information on the physiology and behavior of insects into management programs that are both economically and environmentally sustainable. Currently, much of this research is focused on invasive species, particularly *Halyomorpha halys*, the Brown Marmorated Stink Bug. Anne's disserta-

tion was on the biology of BMSB and was among the first to study this pest in the USA. Her laboratory also evaluates monitoring, biological control, and the integration of management tactics to provide a systems-level approach. This multiple tactic approach is the center of a new \$2.67 million USDA multi-state grant she is leading on BMSB. She looks forward to finding solutions for NJ fruit growers so that we can continue to provide local quality foods.



Dr. Anne Nielsen, RCE Fruit Entomologist, visits a wine grape block at Terhune Orchards, Princeton, NJ with owner Gary Mount. Photo credit: Win Cowgill.

STARK Bro's

A GROWING LEGACY SINCE 1816



Spring planting will be here before you know it, so **now is the time to order.**
Many of our varieties are still available, but they could go fast after the upcoming trade shows.

To guarantee best selection from our crop, call 800-435-8733 today!

Stark Bro's Nurseries & Orchards Co.

PO Box 398 • Louisiana, MO 63353 • 573-754-5111 • Fax 573-754-3701

2013 Mid-Atlantic Fruit & Vegetable Convention
January 29-31, Hershey, PA
(www.mafvc.org)

Tuesday Morning January 29
General Tree Fruit Session

- 9:00 Invocation
- 9:05 President's Address – Carolyn McQuiston
- 9:15 Worker Protection Standards – Jim Harvey, PSU
- 9:45 George Goodling Lecture: Pre and Postharvest Handling of Honeycrisp for Maximum Storage Life and Quality – Chris Watkins, Cornell
- 12:00 Lunch

Tuesday Afternoon January 29
General Tree Fruit Session

- 1:30 The USDA-NRCS partnership with the mid-Atlantic specialty crop industry – Dave Biddinger, PSU
- 2:00 Orchard Renewal Decisions – Clark Seavert, Oregon State
- 2:30 Show & Tell
- 2:45 Vacuum Assisted Harvest for PA Orchard Conditions and Systems – Paul Heinemann and Phil Brown
- 3:30 Comprehensive Automation for Specialty Crops – Looking Ahead after 4 Years of SCRI Funding – Sanjiv Singh, Carnegie Mellon
- 4:15 Adjourn

Wednesday Morning January 30
General Tree Fruit Session

- 9:00 Favorite Cherry Varieties A Grower Panel – Win Cowgill, Rutgers (moderator); Gary Mount, NJ; Evan Milburn, MD; Justin Weaver, PA; Tom Haas, PA
- 9:45 Common Mistakes to Avoid in Training Tall Spindle Trees – Mario Miranda Sazo, Cornell
- 10:15 Show & Tell
- 10:30 Use of SmartFresh on Different Varieties – Chris Watkins, Cornell
- 11:15 New Fire Blight Resistant Pear Cultivars – Richard Bell, USDA
- Noon Lunch

National Peach Council

- 9:00 Late Season Fruit Rots on Peach – Norm Lalancette, Rutgers
- 9:30 Ernie Christ Lecture: A Look at the Past and the Future of Peach Varieties – Jerry Frecon, Rutgers
- 10:15 Show & Tell
- 10:30 Using a phenological model and border sprays for Brown Marmorated Stink Bug - moving towards IPM – Anne Nielsen, Rutgers
- 11:15 Bacterial Spot on Stone Fruit – Sarah Bardsley, PSU
- 12:00 Lunch

Wednesday Afternoon January 30

General Tree Fruit Session

- 1:30 Maintaining Native Pollinator Habitat and Managing Blue Orchard Bees – Dave Biddinger, PSU
- 2:15 Show & Tell
- 2:30 Pesticide Record Keeping Options – Kerry Richards, PSU
- 3:00 US Apple Association Working to Help the Industry – Nancy Foster, US Apple
- 3:30 Pennsylvania Apple Marketing Board Program Update – Karin Rodriguez, PAMB
- 4:00 Adjourn

National Peach Council

- 1:30 National Peach Council Update – Kay Rentzal
- 2:00 Peach Training System Trial – Jim Schupp, PSU
- 2:45 Show & Tell
- 3:00 Peach Rusty Spot Management – Norm Lalancette, Rutgers
- 3:30 Plum and Stone Fruit Hybrids – Jerry Frecon, Rutgers
- 4:00 Adjourn

Thursday Morning January 31

General Tree Fruit Session

- 9:00 YGA on the Move: Young Grower Alliance Panel – Catherine Lara (moderator)
- 9:45 Show & Tell
- 10:00 Pear psylla complex on pears – Peter Shearer, Oregon State
- 10:45 Finding a Good Crop Insurance Program – TBA (Lynn Kime)
- 12:00 Lunch

National Peach Council

- 9:00 Is Your Air Blast Sprayer Calibrated for Your Peach Trees or Apple Trees? – George Hamilton, UNH
- 9:45 Use of Entomopathogenic Nematodes for Borer Control – John Halbrendt, PSU
- 10:15 Show & Tell
- 10:30 Training Young Peach Trees – Rich Marini, PSU
- 11:00 Strategies to Boost Retail Sales – Kathy Kelley, PSU
- 11:30 Grower Experience with Mechanical Blossom Thinning (Grower Panel) – Tara Baugher (moderator); Henry Allenberg, MD; Dave Wenk, PA; Eddie Rankin, PA
- 12:15 Adjourn

Thursday Afternoon January 31

Special Joint Session with PVGA and Small Fruit Program

INVASIVE SPECIES

- 1:30 Brown Mamorated Stink Bug Panel – Tracy Lesky, Greg Krawczyk, Dean Polk
- 2:30 Spotted Wing Drosophila – Peter Shearer, Oregon State, Kathy Demchack, PSU
- 3:15 Invasive Weed Species – Tracy Harpster, PSU
- 3:45 Invasive Vegetable Pests – Shelby Fleisher, PSU
- 4:15 Adjourn

ADVANCED REGISTRATION AND MEMBERSHIP INVOICE
For the Mid-Atlantic Fruit & Vegetable Convention
and Membership to the New Jersey State Horticultural Society
Hershey Lodge and Convention Center – Hershey, PA
January 29, 30 and 31, 2013

Membership Name(s) _____

Farm or Business Name _____

Address _____

City _____ State _____ Zip _____

E-mail Address _____

Telephone _____ Fax _____

Registration Fees

MEMBERS' ADVANCED REGISTRATION # _____ @ \$60 = \$ _____

(Must be postmarked by 1/23/13)

One, Two, or Three Day Members' Registration Will Cost \$75 at the Door

NON-MEMBERS' REGISTRATION (Same for Non Members - Advanced Registration or at the Conf.)

One Day # _____ @ \$100 = \$ _____

Two or Three Days # _____ @ \$140 = \$ _____

GROWER RECEPTION & BANQUET (1/29/2013) # _____ @ \$35 = \$ _____

PROGRAMA de ESPAÑOL (1/30/13) (included in general registration)

PRE CONVENTION PROGRAMS (1/28/13) Register at <http://www.mafvc.org/>

Note: No weather related refunds will be given

CONTRIBUTIONS LECTURE SERIES & RESEARCH

Ernie Christ Distinguished Lecture Series \$ _____

Research Dontations..... \$1000 _____ \$500 _____ \$250 _____ \$100 = _____

Membership Dues

Must be Paid to Receive Membership Rate at Convention @ \$50

UNITED STATES MEMBERSHIP # _____ @ \$50 = \$ _____

(4 Electronic Issues of *Horticultural News*)

Receive a free monthly subscription with your paid membership to NJSHS

_____ *Country Folks Grower* _____ *Fruit Grower News* _____ *American Fruit Grower*

TOTAL ENCLOSED For Registrations, Memberships, and Contributions \$ _____

MAKE CHECKS PAYABLE TO: NJSHS

Return Registration and Dues to:

NEW JERSEY STATE HORTICULTURAL SOCIETY

PO BOX 116, CLAYTON, NEW JERSEY 08312

Web site: <http://www.horticulturalnews.org/>

<http://www.njsHS.org>

Email: polk@njaes.rutgers.edu or frecon@njaes.rutgers.edu before 12/31/12

New Jersey News

Alfred Frank Caggiano Passes

Alfred Frank Caggiano, president of Sunny Slope Farms, passed away peacefully at home Thursday evening, November 8, 2012 with his family by his side.

Sunny Slope Farms grew to be one of the largest peach operations on the east coast. Alfred was a life member of the New Jersey State Horticultural Society and received its Outstanding Grower Award in 1995.

Alfred was born on April 17, 1929 in Elm, NJ to his parents Vincenzo and Antoinette Mangini Caggiano. After graduation, Alfred left home to farm with his older brothers in Bridgeton in the family business. In the 1950's, three of his brothers expanded operations in Gaffney, SC. Alfred, along with his eldest brother, Anthony continued the Bridgeton operation where it grew into a 1,000 acre apple, peach and nectarine orchard. The operation continues today under the guidance of his four children, son-in-law and many faithful employees.

Alfred married Marie Bonnie Halfpenny of Lansdowne, PA. Alfred and Bonnie would have celebrated their 53rd wedding anniversary on November 21st. His involvement with the agriculture industry included service on the NJ Peach Promotion Council, the NJ Farm Bureau and Rutgers' Cook College Association. In addition, Alfred served on the Hopewell Township Planning Board, was chairman of the Cumberland County Board of Health and an original and ardent supporter of West Cumberland

Little League. He also was involved with the formation of the firearm training range for the Bridge Police Department. Alfred was a member of the Board of Directors of Farmers & Merchants National Bank for over 30 years.

Alfred was awarded Bridgeton Area Jaycees' George J. Woodruff Memorial Award, the Bridgeton Area Chamber of Commerce Business Award, and this year was recognized by the NJ Peach Promotion Council who dedicated their Annual Buyers' Guide to him.

Of all his accomplishments, Al was most proud of his family; his wife of more than 50 years, Bonnie; their four children; and seven grandchildren: his daughter,

Lisa and husband Ronald Thomas and their two children; daughter Tara and husband Michael Cintron and son, Tyler; his daughter Cheryl and husband Jeffrey Singer and their three daughters, Kylie, Kendall and Kasey; his son Alfred Jr.; and his daughter Debra and husband Simon Phillips and their two daughters, Allison and Abigail.

In lieu of flowers, memorial contributions can be made in Alfred's memory to Saint Teresa of Avila Church, 46 Central Avenue, Bridgeton, NJ 08302 or to SJ Healthcare-HospiceCare, PO Box 160, Bridgeton, NJ 08302. Written condolences and

tributes may be expressed to the family at the funeral home website, freitagfuneralhome.com



Al Caggiano, Al Caggiano Jr., and Lisa Caggiano Thomas.

Meredith Compton is Named 2013 New Jersey Outstanding Young Farmer

Meredith Compton, a fruit and vegetable farmer and co-owner of Peaceful Valley Orchards in Pittstown, has been named New Jersey's 2013 Outstanding Young Farmer by the State Board of Agriculture and will be honored at the February 2013 New Jersey State Agricultural Convention in Atlantic City.

Compton, 38, also is in the running for one of the top four National Outstanding Young Farmer (OYF) Awards. "Meredith Compton is a dedicated steward of the land who is helping to shape the future for agriculture in New Jersey through active involvement in America's Grow-A-Row, which provides Jersey-grown produce to those in need," said New Jersey Secretary of Agriculture Douglas H. Fisher. "The Outstanding Young Farmer Award helps to recognize young people, like Meredith, for the vital role they play in sustaining New Jersey agriculture." Compton is a first-generation farmer who chose agriculture as

her career after graduating with a Bachelor of Science degree in Agronomy and Environmental Science from Delaware Valley College of Science and Agriculture. She worked at both RAREC and the Rutgers Snyder Research and Extension Farm and then as the North Jersey Tree Fruit IPM program associate for 10 years, honing her agriculture skills. After three years farming on her own while she worked at Rutgers, she decided to pursue farming on full-time basis.

Compton and her husband, Jeremy Compton, operate Peaceful Valley Orchards on leased land. The 150-acre farm offers pick-your-own strawberries, apples, pumpkins and Asian Pears, as well as a Community Supported Agriculture opportunity called the Jersey Fresh Produce Club. Participating members receive a box of fruits and vegetables grown on the farm weekly from July to November.

Peaceful Valley Orchards also is the founding lo-



Jeremy and Meredith Compton

cation of the non-profit organization, America's Grow-A-Row, for which Compton is a board member. For the past eight years, the Compton's have donated land on their farm, as well as seeds and starter plants that are tended by volunteers, harvested and then donated to the hungry.

"We feel that it is essential to provide those in need with fresh fruits and vegetables and to educate people about the important role that agriculture plays in our community and to our well-being," said Compton. "America's Grow-A-Row accomplishes both of those goals and we are proud to say that we have been able to be a part of the success of the program. Grow-A-Row has a lot of exciting plans for the future and we are looking forward to providing many more people in need with Jersey Fresh Produce."

Compton is Vice President of the New Jersey Horticultural Society, a Hunterdon County Board of Agriculture board member, Vice President of the New Jersey Agricultural Experiment Station Board of Managers, and is a graduate of the New Jersey Agricultural Leadership Development Program.


New Jersey annually recognizes the outstanding achievements of a young farmer who is an upstanding leader, respected agriculturalist and is active in community or faith organizations. The Department of Agriculture is now accepting applications for the Year 2014 Outstanding Young Farmer (OYF) Award. The submission deadline for nominations is January 15, 2013. Nominees must be farm operators, deriving a minimum of two-thirds of their income from farming and must be between the ages of 21-40, not becoming 41 prior to January 1, 2014. Candidates who have a financial interest in the farm operation (a sole proprietor, partner, or a corporation) will receive higher ranking

in the OYF judging process. However, nominees may also include salaried farm managers and employees who are actively involved in policy and management decisions. Submission of a financial statement is not required for state competition. A panel of five judges reviews each nominee's application based on the following criteria: progress in their agricultural career (50 percent of the total score); soil and water conservation practices (25 percent of the total score); and contribution to his/her community, state and nation (25 percent of the total score). Any individual or organization can nominate deserving young farmers' names.

Since the first award was presented, eight New Jersey farmers' names have been named National Outstanding Young Farmer. The OYF program is the oldest farmer recognition program in the United States, with the first group of national winners selected in 1955. The goals of the OYF program are to foster better urban-rural relations through the understanding of farmers' challenges, as well as the appreciation of their contributions and achievements; to bring about a greater interest in farmers/ranchers; and to help build an urban awareness of the farmers' importance and impact on America's economy. The National OYF program is sponsored by Deere & Company, supported by the U.S. Junior Chamber of Commerce and the National Association of County Agriculture Agents (NACAA), and is administered by the Outstanding Farmers of America (OFA) Fraternity.

For more information on the state's Outstanding Young Farmer program, visit: www.nj.gov/agriculture/about/sba/cover.html or call Karen Kritz, OYF Program Manager at (609) 984-2506 or email Karen.Kritz@ag.state.nj.us.





Fresher, Sweeter, Juicier

Peaches From Circle M Farms



*Exceptional quality and flavor at attractive prices
Brought to you from the 4th generation family
farm of Santo John Maccherone*



- PLU Stickers Available Upon Request
- Low in Calories
- Nutritious and Delicious
- Exceptional Flavor



SANTO SAYS...

"Peaches from Circle M Farms just plain taste better. Their size and color are what your customers want, and their flavor is the icing on the cake.

That's because our orchards are tended with care and attention, and their fruit is left on the tree to ripen naturally until they reach their optimum sweetness. Then they are picked by hand, taste-tested by me to ensure quality, and packed gently into

25 lb. volume boxes for shipping. Single layer cartons are available upon request.

Circle M's 30 varieties of peaches and nectarines are known for their sweetness and their luscious color. They look as good as they taste.

Discriminating customers appreciate our white flesh peaches because of their lower acid levels and supremely sweet taste.

Add it all up: Peaches from Circle M Farms are fresher and sweeter, picked by hand when their time has come, and packed with care for a longer shelf life.

Yellow, white and donut peaches and yellow and white nectarines are available early July to late September."



**Call us at 856 856 207-4542 or
609 381-6465. Fax your order to
856 878-0404**



Available in 16 oz. and 64 oz. sizes

**Made from fresh New Jersey Peaches
"Peach Cider Drink, Peach Salsa, Peach
Preserves"**

Santo John Maccherone
circlemfarmsllc@gmail.com

**Circle M Farms
88 Sandy Ridge Road
Salem, N.J. 08079**



Horticultural News

New Jersey State Horticultural Society
P.O. Box 116
Clayton, NJ 08312