

Horticultural News

Volume 96, Number 2
Spring, 2016



Horticultural News

Editors: Winfred P. Cowgill, Jr. & Wesley R. Autio

The New Jersey State Horticultural Society was organized on August 17, 1875 at Geological Hall, Rutgers College, New Brunswick, NJ. It remains the oldest Horticultural organization in New Jersey.



Horticultural News began as the *The New Jersey State Horticultural Society News*, in October of 1920. The Society began "collecting paid membership in order to obtain funds to promote new features of the society and extend the usefulness of the society. The Horticultural Society News was started to be the official society publication." Published M. A. Blake, Professor at Rutgers College was the first president and chair of the publication committee.

Editors served as follows:

MA Blake	1920 - 1947
Norman F Childers	1948 - 1980
Win Cowgill	1981 - 1988
Emily Brown Rosen	1988 - 1990
Linda Butenis Vorsa	1991 - 1995
Jerry Frecon	1995 - 2010

June 2010: *Horticultural News* has moved to an online web-based format. The New Jersey State Horticultural Society has partnered with the University of Massachusetts *Fruit Notes*, Dr. Wesley Autio, Editor. Cowgill and Autio will be the new editors of *Horticultural News* and *Fruit Notes*.

Horticultural News is distributed to growers, extension personnel and researchers and libraries across North America. *Horticultural News* focuses primarily on tree-fruit culture, but addresses small-fruit cultural issues as well. Most reports are from current research at Rutgers University, University of Massachusetts, and other universities.

Horticultural News is published four times per year by the New Jersey State Horticultural Society. It is provided as a benefit to membership in the society. Membership costs \$50 per year. Each one-year subscription begins January 1 and ends December 31. Payments via check must be in United States currency and should be payable to the New Jersey State Horticultural Society. *Horticultural News* Electronic subscriptions are available as benefit of membership in the NJ State Horticultural Society. A hidden link will be mailed immediately after publication to members. Issues will be made freely available on this website six months after publication.

Correspondence should be sent to:

New Jersey State Horticultural Society
Greg Donaldson
176 Airport Road
Hackettstown, NJ 07840
TEL (908) 296-1064

For advertising opportunities, contact:

Win Cowgill: wincowgill@mac.com; (908) 489-0207

Table of Contents

Cold Desert Apple Orcharding Kunal Chauhan	1
Discrepancies Between Direct Observation of Apple Scab Ascospore Maturation and Disease Model Forecasts in the 2014 and 2015 Growing Seasons Elizabeth W. Garofalo, Arthur F. Tuttle, Jon M. Clements, and Daniel R. Cooley	7
Growing Peaches in Michigan: How We Do It and What Keeps Us Up at Night Bill Shane	22
Long-time UMass Pomologist and Fruit Notes Editor, Dr. William J. Lord, Passes	24
New Jersey News	26

Cover: Four-year-old apple variety trial trained as a super spindle at Rutgers Snyder Farm, Pittstown, NJ. Win Cowgill photo.

Cold Desert Apple Orcharding

Kunal Chauhan

General Secretary, Progressive Growers Association (PGA), India

Deep in the Himalayas across high mountain passes, lies an extreme place to grow apples. Imagine delicious apples growing at an altitude of 12,139 feet above mean sea level. A place where foreigners would suffer from acute altitude sickness, and the harsh conditions would make life difficult, the locals grow World-class apples out of nothing. Such is the story of the district of Kinnaur and Lahaul Spiti in the Indian state of Himachal Pradesh.

An Overview of the Indian Apple Scene

Apples were introduced to India by the British in the early 1900's, at which time India was ruled by them, the varieties being the classic Bramleys and Pippins. By the time India achieved Independence in 1947, the first orchards had started to flourish in the hills of Himachal Pradesh and in the valleys of Jammu Kashmir. It was the

intervention of Mr. Samuel Evan Stokes, an American living in British India, who felt that the region had the conditions to grow Delicious apples, and he bought five saplings of Starking Delicious from Stark Brothers Nursery. Ever since, the Indian apple market and area under cultivation has grown, with India now being the fifth largest producer of apples in the world.



Figure 2. After the district headquarters of Rekonig Peo, the vegetation becomes sparse, and the orchards are the only speck of green on a high altitude, dry, desert, Leo village.



Figure 1. Orchards on cliffs, the epitome of human endeavour, Pooh village.

Kinnaur and Lahaul Spiti: The Cold Deserts

Apples are grown in the state of Himachal Pradesh mainly in the Shimla district (elevation 4,500 feet to 7,000 feet) where the terrain consists of gentle hills and deep soils. But the place we are talking about is the tribal areas of Kinnaur and Lahaul Spiti district, (elevation 8,000 feet to 12,000 feet), where the best quality apples in India are grown. The climate is dry and winters are cold, and conditions are somewhat similar to Washington



Figure 3. The farms are mostly located in the lower regions of a peak or valley floor and get glacial water from the top of the peaks year-round, which is used as irrigation, Nako village.

State, only the terrain is more rugged. Heading towards Tabo, after the district headquarters of Rekong Peo, the vegetation becomes sparse and the orchards are the only speck of green on a high-altitude, dry, desert.

Climate

Tabo, the place where most of the pictures are from, has a relatively cold environment with the summers seeing highs of around 30 degree Celsius and lows of 10 degree Celsius. Winters are very cold with temperature as low as -13°F. Being a dry desert, the temperature variation of hot days and cool nights in the spring and summer are favourable for the more anthocyanin formation, leading to intense red color and more typey fruit. Almost negligible rainfall means very few diseases and pest.

Practices

The farms are mostly located in the lower regions of a peak or valley floor, and get glacial water from the top of the peaks year-round, which is

used as irrigation. These peaks may have a height of 14,000 to 18,000 feet. Rivers, like Spiti and Satluj, are also present, but that water is not harvested much. Flood Irrigation is mostly practiced by the farmers, with every farmer irrigating the fields once a week.

The only fertilizer they use is farmyard manure and similar. Trees are mostly on seedling rootstocks with Delicious as the main variety and Golden Delicious coming in second. New dwarfing rootstocks are not readily available and are not widely utilized. Most apple trees are large and would remind us of the early 1950's orchards in the US.

The area falls in a rain shadow zone, with minimal rainfall and very low humidity. As a result, most of

these orchard growers do not need to use any fungicides.

The high altitudes of 8,000 feet to 12,000 have cold nights and low minimum temperatures in winter. As a result, these apples have practically zero pests and thus no insecticide usage. Although wooly aphids have been reported from some areas of late, these orchard areas of Himachal Pradesh remain cut off from the world during times of heavy snowfall and will probably remain insect free.



Figure 4. Standard Delicious growing well in Tabo.



Figure 5. The Himalayas run across northwest India and through the state of Himachal Pradesh. The capital Shimla is on the foothills of the Himalayas. The region of Tabo lies on the rain shadow area. The Tibetan Highlands can be seen to the east and the fertile Indian plains to the south.

For the most part, the apples in this region are all natural and hence the fruit has no russetting. The majority of the precipitation is in the winter as snowfall. No plant growth regulators are used, yet the fruit has outstanding typiness. There are minimal temperature variations during the growing season.

Conclusion

The fruit grown in this region is considered to be the best in India. Growers receive a premium price as compared to growers from Shimla area. These Kinnaur region apples retail at \$1.35 per pound in retail stores, as compared to Shimla apples which retail for around \$0.90 for a pound. Washington State

apples retail for \$2.25 per pound in India. Chinese Fuji apples go for \$2.25-\$2.50 per pound. The cost of production of apples for an Indian grower is around 30% of the market price.

Local Folklore

Apart from being an extremely beautiful place and the home for excellent apples, the region also is famous for its various Buddhist shrines, pine nuts, and traditional handicrafts. Located on the proximity of the Tibetan highlands, a lot of influence is found on the architecture of the place. Life is tough at such places, but the place is the epitome of human endeavour.



Kunal Chauhan, General Secretary, Progressive Growers Association (PGA) India. A group of 100+ young farmers (mostly under 35) who have left other careers to pursue the life of an apple grower. His orchard is in the Kotkhai region of Shimla district and mainly grows apples and pears.

FireLine™
OXYTETRACYCLINE

Your solution for -
Fire Blight in Apples & Pears
Bacterial Spot in Peaches & Nectarines



Proven Performers

FireWall™
STREPTOMYCIN

Your solution for -
Fire Blight in Apples & Pear



For the latest on
resistance
management
visit agrosource.net

AGRO SOURCE™

www.agrosource.net

© 2014 AgroSource, Inc. FireLine and FireWall are trademarks of AgroSource, Inc.

PEACHES

*Still growing
strong!*

**Delaware & California Grown
Certified Peach Trees.
Order Now for Spring.**



Adams County Nursery, Inc. • Aspers, PA
(800) 377-3106 • (717) 677-4124 Fax
Website: www.acnursery.com • Email: acn@acnursery.com



Since 1932

The **Best Berry Plants**

- Strawberries, raspberries, blueberries, blackberries, asparagus and more!
- Where the pros go for plans and plants.
- Call for a free catalog and plasticulture guide!

41 River Road, South Deerfield, Massachusetts 01373

NOURSE

www.noursefarms.com 413.665.2658

Proven Performers



Royal Red Honeycrisp®
High color sport of
Honeycrisp. USPP#22,244



Aztec Fuji® DT2 Variety
High color sport of Fuji.
Aztec® Fuji is a protected trademark of
Waimea Variety Management Ltd.



Brookfield Gala®
High color Gala sport with
exceptional shape. USPP#10,016



Lady in Red
High color sport of
Cripps Pink. USPP#18,787



Chelan Spur™
Very compact, high color
red delicious sport. USPPAF

Additional Varieties

Ambrosia™
Blondee™
Cameo® brand
Chrisolyn™ Jonathan
Granny Smith
Jonastar™ Jonagold
Pink Lady® Brand
RubyMac®
Snowsweet®
Spartan
Zestar!®

Gala:

Buckeye® Gala
Ultima Gala®

Early Fuji

Morning Mist™
Rising Sun®

Braeburn

Kumeu Crimson®
Mariri Red™



**Willow
Drive
Nursery**

1-888-548-7337 • 3539 Road 5 NW • Ephrata, WA 98823 • www.willowdrive.com

Discrepancies Between Direct Observation of Apple Scab Ascospore Maturation and Disease Model Forecasts in the 2014 and 2015 Growing Seasons

Elizabeth W. Garofalo, Arthur F. Tuttle, Jon M. Clements, and Daniel R. Cooley
Stockbridge School of Agriculture, University of Massachusetts

Apples are one of the most heavily sprayed crops in the U.S., in part because in temperate climates the disease apple scab (caused by *Venturia inaequalis* (Cke.) Wint) requires several fungicide applications a year for adequate management, and in recent years the number of fungicide applications per season has been increasing in the Northeast (Cooley et al., *Fruit Notes*, Fall 2013). The reasons for this increase are complicated, and include scab resistance to fungicides, the loss of fungicides with significant post-infection activity, and the economic decision that a fungicide application even if scab risk is marginal is inexpensive insurance against significant losses to disease. To reduce fungicide applications, growers need confidence that there is no scab risk. One of the most important tools in estimating scab risk is the model that estimates the maturity of *V. inaequalis* ascospores

The first apple scab infections, primary infections, are initiated by these ascospores, and begin in the spring when ascospores are released from pseudothecia, the fruiting body of the *V. inaequalis* fungus. Pseudothecia develop on infected apple leaves that remain on the orchard floor through the winter (Figures 1 and 2). Once released from the pseudothecia, ascospores can land on susceptible host tissue and, if weather conditions are appropriate, cause infection (Figure 3). These primary infections typically lead to subsequent secondary infections, which can cause scab outbreaks to explode, rapidly spreading through an orchard. While scab epidemics start as primary infections on leaves, it is ultimately secondary infections on fruit that cause the most economic damage (Figure 4). If primary infec-




Figure 1. Upper: apple leaf from an orchard floor showing scab infections from the previous growing season (dark areas). Lower: magnified scab lesion in a leaf showing several fruiting bodies (round, dark objects), pseudothecia, with one pseudothecium circled. (Photos: upper – Ministry of Agriculture, BC, Canada; lower – North Carolina State University Extension)




Figure 2. *Venturia inaequalis* pseudothecium magnified under a microscope (400X) in a prepared "squash mount" showing the three important stages of ascospore maturation: immature asci with no spores or immature spores; a mature ascus with mature ascospores; and an empty ascus which has discharged spores. To estimate maturity, each type of ascus must be counted for several (10) pseudothecia on each evaluation date. (D.R. Cooley)

tions are effectively managed, serious scab epidemics seldom develop.

While there are a number of key elements in effective primary scab management, it is critical that growers know when ascospores first become available, when their numbers are greatest and when their stores have been exhausted. The first release of ascospores marks the beginning of primary scab season, and the last release is the end. Generally ascospore development moves with apple tree development (phenology). Using the cultivar McIntosh as a reference, mature ascospores are generally ready for release when the first green tissue is visible, green tip, and the last mature ascospores are available during early fruit development, a week to 10 days after petal fall.

However, tree development is not always an accurate way to evaluate the chance that ascospores may be released, as tree development varies by cultivar. Additionally, in some years weather may delay or advance ascospore development relative to tree growth stage. If ascospore release starts after green tip, growers will spray fungicides unnecessarily early in the season. If the primary season lasts longer than estimated, growers may not apply fungicide against late-season primary infection events, and have to battle scab for the rest of the growing season. The early season, around green-tip, is often considered relatively low risk in "clean" orchards, because the relative amount of scab inoculum is low. However, if ascospores development is advanced, the relative amount of inoculum increases, raising risk of

infection.

A much more accurate way to determine whether ascospores are mature and ready to be released, is to look for them. Because ascospores cannot be observed by the naked eye, it takes a microscope and a trained observer to run such evaluations. In the early days of apple IPM, Extension made direct ascospore maturity observations using microscopes in laboratories, a process that involved significant time and some skill.

To eliminate this labor-intensive process, researchers developed models that relate ascospore maturity to growing degree-day (GDD) accumulations, a means of measuring heating units, often calculated by determin-



Figure 3. Primary scab infections on an apple leaf. (Photo APSnet.org)



Figure 4. Apple scab on an apple that was not sprayed with fungicides. (D.R. Cooley)

ing the mean of the high and low temperature for the day (James & Sutton, 1982; Gadoury & MacHardy 1982; Figure 5). One of these developed at the University of New Hampshire has been widely used in IPM programs. GDDs are calculated using a base of 32°F, and accumulation is started at 50% green tip of McIntosh. The model enables users to estimate periods when a large proportion of ascospores are available, and the end of primary scab season, with somewhat better accuracy than tree growth stages. In a typical year, these times could be predicted once green tip occurred, based on historical temperature data (MacHardy & Gadoury, 1985).

It was noticed, however, that ascospore maturation could deviate from the model, sometimes significantly. While the New Hampshire model used only temperature, a model developed in North Carolina had also included a moisture factor, requiring rain or high relative humidity for ascospore maturation. Based on observations of scab infections in Norway, researchers re-examined the New Hampshire model to see whether a moisture factor would improve estimates of ascospore maturity (Stensvand et al, 2005). They found that as little as four consecutive dry days, with less than 0.2 mm of precipitation and less than 12 hours of dew or




Figure 5. Graph showing the maturation of *Venturia inaequalis* ascospores in response to degree-days, base 32 °F accumulated from green tip. The narrower lines are 90% confidence intervals indicating the range of ascospore maturation at a given number of degree-days 90% of the time.

fog, is enough to cause ascospore maturity to slow, and if these dry conditions continued for 7 days or more, maturation ceased altogether until wetting requirements have once more occurred. This phenomenon is technically called a rainfall frequency threshold, and commonly referred to as the “dry switch”. This led to an adjustment to the model to deal with dry periods

The New Hampshire ascospore maturity model has been incorporated into web-based Decision Support Systems (DSSs), which combine weather data with pest forecasting models to provide online advice to growers about pest risks and treatment options. There are different DSSs available in New England and New Jersey to estimate risk of scab, all of which use variations of the MacHardy/Gadoury ascospore maturity model. These include NEWA (the Network for Environment and Weather Applications) managed by the New York State IPM Program at Cornell; Ag-Radar managed by the University of Maine Extension; and the commercial product SkyBit (ZedX, Inc.). Additionally, RIMpro, developed and maintained by Bio Fruit Advies in the Netherlands will be available on a limited, trial basis for growers in northeast in 2016. In a typical growing season, the models in these DSSs are accurate in New

England. However, when growing conditions stray from those generally experienced, failures occur. Indications are that more extreme weather events, such as prolonged dry or wet periods, as well as increasing temperatures, now occur and will continue (Kunkel et. al. 2013). This was clearly seen in 2015 when drought came to the Northeast in the spring, a typically wet time of year. The National Drought Mitigation Center’s US Drought Monitor reports that 23% of the Northeast experienced a moderate drought in May, and that up to 64% of the Northeast experienced abnormally dry weather thorough April and May. The number of frost-free days is also increasing. Subsequently, timing of green-tip has become more variable, making it an even less appropriate indicator for the presence of mature ascospores.

The difference between DSS estimates and actual ascospore maturity was seen in 2012 in the Hudson Valley of New York, where the NEWA DSS estimated a much earlier end of primary ascospore production than was observed using a laboratory evaluation of ascospore release from leaves (Rosenberger, personal communication). While NEWA uses a dry switch, it apparently is not adequate for a very extended dry period with less rain and lower relative humidity levels than had been seen in that area since ascospore maturation levels had begun to be documented. Subsequent discussions with a group of researchers suggested that thresholds for “dry” and “wet” in the ascospore maturity model needed to be improved.

In an attempt to better understand how weather such as 2012 might affect the New Hampshire ascospore maturity model and DSS estimates of scab risk, in 2014 and 2015 we compared DSS estimates of ascospore maturation with different types of direct ascospore maturity observations. In the fall, scab-infected leaves from unsprayed apple trees were collected and overwintered outdoors at the University of Massachusetts Cold Spring Orchard Research and Education Center (CSOREC) in Belchertown, MA. In the spring, these leaves were used in direct observations




Figure 6. Petri dish with microscope slides on the bottom section, and overwintered, scab-infected apple leaves attached to the cover section. (L. Garofalo)

of ascospore maturity. In 2014, we used three assays:

1. Standard “squash mounts” of fruiting bodies of the fungus sampled from a set of leaves (Figure 2);
2. A Petri-plate assay in which, after wetting, leaf sections are put in a moist Petri plate and spores are released onto a microscope slide (Figure 6);
3. A spore trap using fans to concentrate on a microscope slide ascospores released from several leaves, the “two-fan trap” (Figure 7).
4. In 2015, we added a fourth method: A spore trap in the lab using a single fan to draw air over leaves down through a funnel to an orifice over a microscope slide, the “funnel trap” (Figure 8).

For the squash mounts, 10 pseudothecia were collected from six randomly selected leaves. The pseudothecia were placed in a drop of water on a glass microscope slide, and gentle pressure applied to a glass cover slip on top of them to break them open. The asci and any ascospores inside the pseudothecia were then evaluated for maturity.

Petri-plate assays were performed by tapping six sections of overwintered leaves containing pseudothecia into the lid of a Petri plate. Two microscope slides were placed in the bottom of the Petri plate with a small amount of distilled water, and the lid placed on top. After an hour, these slides were examined for the presence of released ascospores. These lab tests were performed approximately every week.

Ten to fifteen leaves with pseudothecia were used for the field-based spore trap. These were placed in a single layer on the ground under a large-mesh wire screen. The leaves were surrounded by a rectangular “box” with dimensions of 15 ¾” X 9 ¾” in. X 8”, framed with wood strips, open sides were covered with landscape fabric and the ends were of hardboard. Within the box leaves were held in place,

scabby side up, with hardware cloth. Two fans were located at opposite ends of the box, which were activated by a leaf wetness sensor when a rain event began. The fans moved an air current from one end of the box to the other, where any released spores were forced through a small opening onto a microscope slide. The slide was brought back to the lab and inspected after each rain event ended.

The funnel trap was used on leaves brought from the field to the laboratory. It consisted of a plastic funnel with a small electric fan that drew air over wet leaves placed on a screen at the wide end of the funnel down to a slot and microscope slide at the narrow end of the funnel. The fan was run for 20 minutes, after which the



Figure 7. Two-fan trap designed by MacHardy, actual trap with leaves on the left, schematic showing design on the right. (W.E. MacHardy)

slide was inspected for spores.

We compared four DSSs: NEWA, Ag-Radar, SkyBit, and a commercial product being introduced in the U.S., RIMpro (Bio Fruit Advies, Zoelmand, Netherlands). The DSSs have rules that determine when maturation will start (the biofix), usually a simple observation of green tip as in the New Hampshire model. SkyBit estimates a green tip date, which can be corrected by supplying ZedX with the observed green tip date. RIMpro uses the first date that spores are released, or green tip, whichever comes first, as the biofix.

Weather data for the NEWA maturity estimates came from a weather station (Rainwise, Trenton, ME) at the CSOREC site. RIMpro also used data from this station. SkyBit uses virtual weather data. Virtual data are created by combining different sources of actual weather observations (e.g. National Weather Service) with proprietary mathematical techniques, which basically interpolate from the actual observations to estimate weather for locations distant from weather stations. SkyBit sells E-Weather service products and offers an “Ag-Weather IPM Apple Disease Product” that includes virtual weather

data and scab risk. Ag-Radar uses the SkyBit virtual data in a somewhat different model.

In 2014, ascospore observations were not made until green tip on April 14, at which point mature ascospores were observed in all three assays (Table 1).






Figure 8. Funnel trap designed by MacHardy, showing leaves in wire mesh. Air is drawn down to an orifice over a microscope slide. (W.E. MacHardy)



The Petri-dish assay and the field trap had similar patterns of ascospore maturation, while the squash mount assay had a maturation pattern that lagged behind the trapping assay (Figure 9). The direct observations of last available ascospores, indicating the end of primary seasons, were very different. For the Petri-dish assay, the last spores were seen on May 19, for the two-fan field trap on June 2, and for the squash mounts, June 9.

In 2014, the four DSSs also differed in terms of their estimates of the end of primary apple scab (Table 1). NEWA had the earliest estimate of the end of ascospore release, May 28. RIMpro, SkyBit and Ag-Radar estimated the end of the season within 3 days of each other, on June 3, June 4 and June 5, respectively. That pattern of maturation was generally similar for the DSSs, though SkyBit was consistently the latest and RIMpro the earliest (Figure 10).

In 2015, squash mounts were the first to show mature ascospores on April 6, two weeks before McIntosh green tip (Figure 11). Though the spores appeared mature in the squash mounts, no spores were released in any of the trapping assays until three weeks later on April 27 (between green tip and tight cluster), when spores were seen in the Petri plate assay. First release in the funnel trap assay and field two-fan trap were even later, at May 4 (tight cluster) and May 19 (petal fall), respectively, one to three weeks later than the first spores seen using the Petri-plate trap. Overall there was a difference of nearly six weeks in first mature spore observations between the squash mounts and the last trapping assay, the field trap (Table 2).

Similarly, squash mounts estimated the earliest ascospore development, though the Petri dish and funnel trap assays were similar to squash mounts after petal fall (Figure 11). The two-fan field trap indicated the slowest spore development; no spores were caught in it until after petal fall. The assay estimates of the end of primary season were more consistent. All were on the same day, June 22. (Table 2)

The DSS estimates of the beginning of scab season were more consistent among each other. NEWA and Ag-Radar estimated the same day, April 19 (green tip), while RIMpro and SkyBit estimated later dates, April 22 and April 27, respectively. The DSSs differed by over a period of over three weeks in their estimates of the end of primary season, from May 27 by NEWA to June 20 by RIMpro (Table 2). RIMpro had an extended period of no maturation, 22 days from April 29 to May 20. Ag-Radar showed no spore development from April 30 to May 12 (13 days), NEWA from April 29 to May



4 (6 days), and SkyBit from April 27 to May 3 (7 days) (Figure 12).



These periods reflect a prolonged period of dry weather that began on April 23 and ended May 19, when 0.13 inches of rain was recorded by the weather station. The weather station also recorded 0.02 inches of rain, the minimum recordable amount, on May 5 and again on May 12. (Figure 14). SkyBit estimated 0.02 inches of rain on May 12, and 0.42 inches on May 19.

In 2015, NEWA, SkyBit and Ag-Radar estimated the end of primary season to be two to three weeks earlier than the last date when ascospores were actually observed. The RIMpro estimate, June 20, was much closer to the last spore captures, June 22. It is clear that the different systems have different ways of estimating the impact of dry periods on spore maturation. The assays all indicated that mature ascospores were available and released from two to four weeks after petal fall, much later than the usual week to 10 days. Both NEWA and SkyBit estimated the last ascospores were available at the end of May, two to three weeks before the last spores were seen in the assays. Ag-Radar estimated June 8 as the end of ascospore production, one to two weeks earlier than the assays. RIMpro estimated an end of ascospore production to be June 20, only two days off from the June 22 assay date.

Both the DSS estimates and the assay observations matched tree phenology in 2014, a normally wet year without extended dry periods (Figure 13). In 2015, with the extended dry periods, assay observations and estimates from three of the DSSs were significantly different. From a management perspective, this means that growers depending on NEWA, SkyBit or Ag-Radar to determine the end of primary inoculum availability for apple scab would stop scab fungicide applications prematurely. For example, NEWA states “Ascospores were essentially all released on May 27” and that infections from that event should appear within 10 days by June 6. Four infection periods occurred between June 6 and June 22, when the last ascospores were trapped. If fungicides were not applied to protect against those four infection periods, there would have been a risk of primary infection. RIMpro was the one DSS that accurately estimated ascospore maturity as it was observed.

An important aspect of DSS recommendations is using weather forecasts to predict the risk of scab infections. Since fungicides should be applied as protectant materials, before infections happen, growers need to know when scab infections are likely to occur. Such forecasts use models which are based on Mills





infection periods with different modifications, and the accuracy of the predicted risk of infection depends both on the model and the weather forecast. We do not present the details here, but generally DSSs predicted potential wetting and infection periods a few times at

CSO during May, but they never developed. Growers could have decided to wait to see whether a forecast infection period in fact occurred, and then have applied a post-infection fungicide, but if preventive fungicides were applied before the predicted infection, they were

Table 1. Comparison of ascospore maturation in terms the end of primary scab at Cold Spring Orchard, Belchertown, MA, 2014

Spore Assay Method	First Mature Ascospores*	Over 95% of Ascospores Mature	Last Observed Ascospores
Squash mount	April 14	June 9	June 9
Petri plate	April 14	May 16	May 19
Two-fan trap, field	April 14	May 17	June 2
Decision Support System			
NEWA	April 14	May 20	May 28
AgRadar	April 14	May 23	June 5
RIMpro	April 14	May 21	June 3
Skybit	April 14	May 26	June 4
Tree Phenology McIntosh			
Green tip	April 14		
Petal fall plus 14 days			June 2

*First ascospore assays were performed at green tip on April 14, when all assays showed mature ascospores. DSS estimates of first spores are made prior to supplying actual green tip dates.

Table 2. Comparison of ascospore maturation in terms of first date (beginning of primary scab) and last date (end of primary scab) at Cold Spring Orchard, Belchertown, MA, 2015

Spore Assay Method	First Mature Ascospores	Over 95% of Ascospores Mature	Last Observed Ascospores
Squash mount	April 6	June 1	June 1
Petri plate	April 27	June 8	June 22
Two-fan trap, field	May 19	June 15	June 22
Funnel trap, lab	May 4	June 8	June 22
Decision Support System			
NEWA	April 19	May 22	May 27
AgRadar	April 19	June 2	June 8
RIMpro	April 22	June 14	June 20
Skybit	April 27	May 25	May 31
Tree Phenology McIntosh			
Green tip	April 19		
Petal fall plus 14 days			June 2

unnecessary. In the worst case, a grower relying on DSS information might have applied two to three unnecessary fungicides during May, and not applied fungicides when they were still needed in June.

Clearly, further modifications of some DSS models are needed in order to better estimate when fungicide applications are needed and when they are not. Information from the DSSs is useful, but needs to be tempered by grower experience and timely information from other

sources. For example, during prolonged dry weather in primary scab season, growers could plan on using post-infection tactics so as not to over-apply fungicides, and be prepared to apply scab fungicides after fruit set in case observations indicate primary scab season extends beyond that point. In time, model accuracy and weather forecasts will improve, but grower experience will undoubtedly always play an important role in efficient and effective scab management.

Literature Cited

- Clements, J. "A New England-wide Apple Decision Support System For Apple Scab, Fire Blight and Coddling Moth." 77th Annual New England, New York and Canadian Fruit Pest Management Workshop. Booth's Bishop Conference Center, Burlington, VT. 20 October 2015. Research Presentation.
- Cooley, D.R., A. Tuttle, S. Villani, K. Cox, G. Koehler, T. Green and P. Werts. 2013. Increasing fungicide use in New England apples. *Fruit Notes* 78 (4) 1-6.
- Gadoury, D. M. & MacHardy, W. E. 1982. A model to estimate the maturity of ascospores of *Venturia inaequalis*. *Phytopathology*, 72, 901-904.
- Gadoury, D.M., Seem, R.C., MacHardy, W. E., Wilcox, W. F., Rosenberger, D. A. and Stensvand, A. 2004. A comparison of methods used to estimate the maturity and release of ascospores of *Venturia inaequalis*. *Plant Disease*, 88: 869-874.
- James, J. R. & Sutton, T. B. 1982. A model for predicting ascospore maturation of *Venturia inaequalis*. *Phytopathology*, 72, 1081-1085.
- Kunkel, K. E., L. E. Stevens, S. E. Stevens, L. Sun, E. Janssen, D. Wuebbles, J. Rennells, A. DeGaetano, and J. G. Dobson, 2013: Regional Climate Trends and Scenarios for the U.S. National Climate Assessment: Part 1. Climate of the Northeast U.S. NOAA Technical Report NESDIS 142-1. 87 pp., National Oceanic and Atmospheric Administration, National Environmental Satellite, Data, and Information Service, Washington, D.C. [Available online at http://www.nesdis.noaa.gov/technical_reports/NOAA_NESDIS_Tech_Report_142-1-Climate_of_the_Northeast_U.S.pdf]
- McHardy, W.E. and Gadoury, D.M. 1985. Forecasting seasonal maturation of ascospores of *Venturia inaequalis*. *Phytopathology* 75: 381-385.
- Stensvand, A., Eikemo H., Gadoury, D.M. and Robert C. Seem. 2005. Use of a rainfall frequency threshold to adjust a degree-day model of ascospore maturity of *Venturia inaequalis*. *Plant Disease* 89:198-202.
- National Drought Mitigation Center. (2016). [Table of dates describing various stages of drought and percent of region experiencing them. Regional map]. United States Drought Monitor. Retrieved from <http://droughtmonitor.unl.edu/MapsAndData/MapArchive.aspx>.

Eco-Friendly Insect, Disease, Bird Control

University/USDA tested

Stink Bug Traps

Brown Marmorated and Native Bugs

Insect Traps and Lures

*Plum Curculio Trap Tree Control,
Codling & Oriental Moth, Cranberry
Pests, Many Others*

Honey Bee Lure

Attract Bees - Increase Pollination

Oriental Beetle MD

*Mating Disruption
Fruit Crops & Ornamentals*

Avex

*Bird, Goose, Duck Repellent
Berries, Fruits, Row Crops*

Agri-Fos

*Phytophthora, Pythium
The "original" phosphite fungicide*

AgBio Inc.

*Committed to the Environment and Green Technology
Since 1990*

P. 877-268-2020
agbio@agbio-inc.com
www.AgBio-Inc.com





SUMMIT TREE SALES

REPRESENTING THE COUNTRY'S FINEST NURSERIES

Make Summit your first call!

Your time is valuable. Let the highly qualified staff at Summit find the trees you need. Summit Tree Sales has a combined 50+ years of experience in the fruit industry.



Jan Melvin



Katie Schuld

Let Summit manage *all* your orders from start to finish!

When you book through Summit, you will have access to your current and past orders with a simple phone call. Summit will book all your contract and non-contract orders, make any requested order changes, and work closely with the nursery to make sure you receive the trees you need.



Dawn Melvin



Matt Schuld

Our Service is Free!

Your price is the same as when ordering directly from the nursery.

Don't be left out - Order Trees Today!

800.424.2765
summittreesales.com

***THE COMMERCIAL GROWER'S
ONE-STOP SOURCE FOR FRUIT TREES***

SUMMIT TREE SALES | 800.424.2765 | SUMMITTREESALES.COM
55826 60TH AVE LAWRENCE, MI 49064



Quality Ag Equipment



Built to Last Since 1986

COMMERCIAL ORCHARD & PRODUCE EQUIPMENT



Model	Length	Width	Bin Capacity	Weight Capacity
12-1022DOBT	22'	120"	12	24,000#

DECKOVER FIELD PACKING WAGONS

- Conveyor Compatible
- 2 Rows of Bins with Center Aisle
- 4# Steel Grate Decking
- Walking Beam Axles Standard

LOW-PRO BIN TRAILERS

- Grip Strut Steps for Safety
- 4# Steel Grate Decking
- Walking Beam Axles Standard
- GAP Required Accessories



Model	Length	Width	Bin Capacity	Weight Capacity
8-524LPBT	24'	60"	7	8,000#



HAYRIDE WAGONS

- Built to Suit Customer Needs
- Industry Leading Safety Features
- Multiple Seating Configurations
- Folding & Stationary Steps
- Roof Options Available

Optional Walking Beam Axles



Optional Hitches



Optional LED Light Packages

Covered by
Diller's Exclusive
**3 Year
Warranty**
See Warranty Guide for
Coverage Details

Factory Direct Sales to NJ & MA
800-523-8002 or visit us at: www.DillerAg.com



2012 Diller Ag Equipment • a Division of eXtreme Manufacturing

**It's worth the time to review your weather risks,
when you think of the damage caused in just a flash.**



A Whole-Farm Revenue Protection policy working along with other crop insurance policies is a great way to minimize your revenue risk. Protect your income from whatever Mother Nature has in store.

Contact us today to learn more at **877-867-9291**.

11539 Nuckols Road, Suite B; Glen Allen, VA 23059

**AgRISK
MANAGEMENT**

By Weller & Associates, Inc.

agRISKmgt.com

Harvest time and the pickin' is easy



REVO

The REVO Piuma 4WD Harvester

- For apple & pear high-density orchards
- Picking, pruning & trellis work
- Independent front & rear steering & crabbing for tight turns
- 12'-6" footprint
- Automatic self-leveling system
- Whisper-quiet diesel engine
- Onboard compressor for air-driven tools
- Compact / no trailer to pull
- Flow-thru bin design

**Call us for a demonstration
800-634-5557**

Scan this code
to see the
Piuma in action



60
YEARS

Supplying Growers, Gardeners

OESCO, INC.
and Groundskeeping Professionals. Since 1954.

P.O. Box 540
Rt. 116 / 8 Ashfield Road
Conway, MA 01341
www.oescinc.com



Growing Peaches in Michigan: How We Do It and What Keeps Us Up at Night

Bill Shane

Michigan State University

Making money producing peaches is difficult because the tree is sensitive to low winter temperatures, blooms relatively early, and the fruit has a short storage life. However, selling good quality peaches is generally easy because of the demand for this queen of fruit. Farm marketers know that peaches draw customers to their fruit stand. I am providing here observations and strategies that I have learned over the years working in Michigan for increasing the productivity of peaches. Although these tips will not eliminate the anxieties associated with growing peaches, I am confident that some of these will help.

Tree Longevity Anxiety

Temperatures below approximately -13 F in the mid-winter is tough on any peach tree and certainly on peach fruit buds. There are what I call lower tier varieties such as Veteran, Reliance, and Madison, that have the reputation for the best mid-winter hardiness, but their fruit quality and/or appearance are only so-so. 2nd tier peach varieties not quite as hardy as these but with better quality include Harrow Diamond, Starfire, Contender and the flat white fleshed peach Saturn. A third tier of varieties slightly less hardy reliable than these, but still pretty good, and good to excellent quality include Garnet Beauty, Summer Serenade, Redhaven, Allstar, PF17, and Glowingstar.

Since winter damage and tree mortality is a fact of life with peaches, a good strategy is to have an ongoing orchard planting strategy to insure a farm has a range of trees ages, and at least two varieties for each harvest window. A farm with a range of tree ages will have a better chance of surviving a cold winter with some blocks still viable. Winter damage tends to be worse on older trees, but not always.

A careful consideration about the planting site is important to help head off problems. Peach tree do best on

sandy loam soils with 3 or more feet thick of topsoil. Tile drainage systems are critical in many orchard sites to handle excess water. In addition to tiles, soils that have more silt or clay than ideal should be shaped into a slight berm so that excess water will drain out of the root zone area. Although peaches do better on sandy sites than other tree fruit, trickle irrigation is invaluable to get an orchard started in good shape during droughty years.

Tree planting depth is particularly important on heavier soils. The traditional approach is to plant peaches so that the graft union is at the soil line. How-



Figure 1. Bill Shane, peach tree specialist at Michigan State University SW Research Center.

ever, if the shank (the part between the graft union and the topmost root) is long (greater than 1 foot), the tree should be planted so that the topmost root is within a few inches of the soil line, which will put the graft union above the soil line. Tree roots that are planted too deep in heavier soils are prone to collar and root rot problems.

For sandy sites a more subtle concern is potential tomato ringspot virus problems. Peach trees with this virus are more prone to reduced productivity and shorter lifespan. This virus is spread from weed to tree and tree to tree by the dagger nematode, a pathogen that prefers sandy sites. The only two reliable ways to check a site for this problem are to test for the dagger nematode or to test the known weed hosts such as dandelion or the tiny root of the peach tree for the virus. Both tests can only be done by labs with the proper diagnostic kits. Growers who do a good job of keeping broadleaf weeds out of their orchards generally have less problems with this virus. Sites with this virus and nematodes are good candidates for pre plant fumigation or other nematode-fighting techniques such as use of rapeseed, mustards, or other non-host rotations.

Another potential problem for us in Michigan is X-disease, caused by a phytoplasma, which is somewhat like a bacterium, but without a cell wall. Trees with X-disease will develop a characteristic red wine colored leaf spot, drop their older leaves, and then decline and die within a few years of infection. This disease is spread by several species of leafhopper, which explains why the disease appears sporadically. The other clue to disease spread and control is the fact that leafhoppers acquire the pathogen from infected chokecherry, tart cherry, and sweet cherry. When X-disease starts showing up in a peach orchard, it usually means that the grower needs to go on a witch hunt for the chokecherries or possibly old tart or sweet cherry trees that are serving as the source of X-disease that the leafhoppers are acquiring—a source that could be a ¼ mile or more away.

Nursery Tree Anxiety

Nursery tree quality has a big effect on the productivity and useful

lifespan of a peach orchard, but is sometimes out of the control of the grower. I have seen cases where nursery trees were exposed to ethylene from apples stored in the same building and the trees would not grow properly. Another occasionally seen problem is trees that grow properly for a few years, but then show trunk splitting and root suckering because the trees had trunk cambium damage due to cold while in the nursery. Diagnosing this requires dissecting a few trees to look for dead cambium tissue in the inner rings of the tree. Another difficult to diagnose problem is trees that have roots dried out somewhere between digging in the nursery and planting. Such trees will be slow to grow in the new planting. I generally recommend that growers plant trees from more than one nursery in a new orchard so that these types of problems are more easily diagnosed.

Tree quality can be particularly important for the success of certain training systems. For example, the Y and quad training systems requires that the nursery tree be headed low at the time of planting. A low vigor tree that pushes out little growth after this heading cut will offer poor options when it comes time to select branches for the scaffold arms. The effort and time spent hunting for the right type of tree is often well spent. I have seen some growers in Michigan preferring June-budded peach trees produced by some nurseries because of their smaller caliper and greater readiness to grow when headed low.



Figure 2. Southwest Michigan is a major producer of high-quality, fresh-market peaches.

Training System Anxiety

Some growers have the knack for growing long-lived orchards. These growers tend to take the time needed for site preparation techniques mentioned above. In addition, they also use tree training techniques that encourage good tree structure. There are many training systems such as 3 to 5 scaffold open center, central leader, Y, palmette, fusetto, quad, and many variations within any one system. The key feature of a long-lived tree is that the scaffold limb arrangement avoids “plumbing” problems. One way to visualize this is to think of a tree as a plumbing project, with tubes (xylem and phloem) running just under the bark. A well-structured tree provides relatively unimpeded flow between the trunk and the scaffolds. A scaffold limb that is “stacked” directly above another scaffold has no clear access to water flow from the roots. Two scaffold limbs that are side by side block “flow” to limbs above them.

Some growers rely on old fashioned wooden clothes pins to help insure good plumbing in their trees. The trees are clothes-pinned when potential new scaffold limbs are 4 to 6 inches long. The clothespin is clamped on the central leader above the new limb such that the tails of the clothespin direct the limb to grow horizontally. This helps to avoid bad crotch angles, poor limb strength, and poor circulation.

Another trick for early years of an orchard is to use 2 or three rounds of pinching and limb breaking in the spring to early summer to encourage growth elsewhere on the tree. This is the so-called “benign neglect” approach to training, a term coined by University of California Extension Specialist Kevin Day. The presence of the broken limbs helps to prevent strong regrowth which happens if the limbs are simply pruned. The tree “gives up” on the broken limbs which are eventually pruned out, but the impact is less harsh than making strong cuts on a young tree.

A third technique to avoid problems in a young tree is to insure that the scaffold ends remain simple in the 2 or 3 years of growth. At the time of bud swell, the excess buds in the first 4 or 5 inches at the scaffold end are removed, leaving the end bud or two intact. This helps eliminate the need for later strong cuts to remove the excess cluster of limbs, a harsh pruning that often leads to disease canker problems. Debudded trees need to have a good spray program for oriental fruit moth to protect the few remaining buds on the scaffold ends.

Bill Shane is a Senior Tree Fruit Extension Specialist and Peach Breeder at Michigan State University. He is also the new director of the South West Michigan Research and Extension Center located in Benton Harbor, Michigan

Long-time UMass Pomologist and Fruit Notes Editor, Dr. William J. Lord, Passes



It is with great sadness that we report the passing of Dr. William J. Lord. Bill served at UMass as the Extension Tree Fruit Specialist, Professor of Pomology, and Editor of Fruit Notes from 1955 to 1985. Bill passed away on March 19 at age 94.

After retirement, Bill focused on fly fishing, but he continued to teach and work at the UMass Cold Spring Orchard for many years. His enthusiasm for pomology and for the orchards of Massachusetts never waned.

Donations in memory of Bill should be directed to the Winifred C. and Jesse L. Rice Fund, UMass Cold Spring Orchard Research & Education Center, 391 Sabin Street, Belcher-town. The income from this fund supports the operation of the UMass Cold Spring Orchard.

The Rootstock Bottleneck Stops Here!



APPLE
CHERRY
PEAR
PEACH
ALMOND
WINE GRAPE
MORE!

**WE CAN WORK WITH
YOUR NURSERY!**

WE ALSO OFFER THESE SERVICES:

- DNA Based Identification -
Genetic certification
- Plant Repository Service
- Virus Screening

www.phytelligence.com
206-719-5317

New Jersey News

Ernie Christ Memorial Lecture, 2016 Mid Atlantic Fruit and Vegetable Convention

Win Cowgill

New Jersey Agricultural Experiment Station

The Ernie Christ Memorial Lecture is presented at the Mid Atlantic Fruit and Vegetable Convention in February each year, held in Hershey, PA. The lecture was established in memory of Ernie Christ, the long time New Jersey Tree Fruit Specialist at Rutgers Cooperative Extension. Ernie passed on September 12, 2000. He was loved and respected by fruit growers across North America. Ernie's passion was the furthering of knowledge of peach culture and science.

Ernie was a great friend and mentor to dozens of young scientists, extension workers, and farmers, including Rich Marini, Sue Brown, Jerry Frecon, Mark Robson, Robert Best, Peter Melick, Ken Wightman, Bob Schutzki, Al Caggiano, Ralph DelSanti, Jack Springer, Spence Davis, John Gerwig, Gary Mount, Bill Smith, Galetta Farms, Santo Maccherone, Spitz Tomasello, and me, to name only a very few! We referred to him as Mr. Peach. The fruit industry thought so much of Ernie they named a NJAES breeding program selection after him, 'Ernie's Choice' which is still grown today.

A speakers fund was established by the NJ State Horticultural Society with an initial gift by Adams County Nursery, with continued funding from grower donations.

The fund supports an invited speaker each year at the Mid Atlantic Conference. The first Ernie Christ Memorial Lecture was presented by Dr. Rich Marini, Horticulture Department Head, Penn State University, in January of 2002.

Professor Dean Polk, Rutgers, presented the 2016 annual Ernie Christ Memorial Lecture, titled "Getting Back into an IPM Program in Peaches & Nectarines." Meredith Compton, President, New Jersey State Horticultural Society, presented Professor Polk with a certificate and a cash award for his invited lecture.



Two Awarded Distinguished Service to New Jersey Agriculture Citations

Lynn Richmond

NJ Department of Agriculture

In Atlantic City, New Jersey at the NJ Agricultural Convention, January, 2016, Rutgers Cooperative Extension retirees Drs. Gerald M. Ghidui of Bridgeton and Bradley Majek of Pittsgrove, were honored February 10 with Distinguished Service to Agriculture Citations by the New Jersey State Board of Agriculture.

Gerald Ghidui

“New Jersey is fortunate to have such outstanding individuals who have dedicated their lives to the betterment of the Garden State’s agriculture industry,” said New Jersey Secretary of Agriculture Douglas H. Fisher. “The state’s farmers owe a lot to Dr. Ghidui and Dr. Majek. Their contributions have had a lasting impact on farmers and young people interested in agriculture.”

Dr. Gerald Ghidui’s service to New Jersey agriculture began in 1980 when he became a Rutgers University extension Entomologist. His career spanned 33 years during which he worked with farmers and conducted research to assist them in controlling vegetable pests.

Dr. Ghidui earned a BS in Biology and Secondary Education from the State University of New York and a PHD in Economic Entomology from Iowa State University. He then worked for the USDA Federal Corn Insects Research Unit in Iowa, focusing on the European corn borer.

While at Rutgers, he presented many educational sessions to students, growers, ag specialists and agribusiness personnel. His research centered on the Colorado potato beetle, European corn borer, and insect pest management of vegetable insect pests. He developed many practices that are now commonplace

in the industry.

Dr. Ghidui gave presentations and lectured throughout the country on his work and was published in numerous journals, papers, magazines and newsletters.

From 1996 to 2000, he was the Director of the Rutgers Agricultural Research and Extension Center in Bridgeton. He served as Supervisor of the IR-4 Regional Research Center at RAREC and was the IR-4 State liaison. IR-4 activity helps connect growers, the agrichemical industry and federal agencies to secure clearance for pest control products on vegetables, eventually becoming a registered label for use on that crop. He completed a total of 84 IR-4 projects for insecticides over his career. He was editor of the Journal of Economic Entomology, Entomological Society of America. And he served as Editor of the insect control section of the NJ Commercial Vegetable Production Recommendations. Dr. Ghidui retired Professor Emeritus from Rutgers in September 2012.



Secretary Fisher awards Gerald Ghidui with Lilly Ghidui the Distinguished Service Award from the NJ Department of Agriculture.

Bradley Majek

Dr. Bradley Majek has played an important role in New Jersey agriculture for more than 34 years. He worked at Rutgers University conducting research and extension programs in tree fruit and vegetable crops since 1981 and has filled a critical need of improving crop productivity in New Jersey. Majek earned his BS from Cornell University, his Masters in Weed Science and Botany from Oregon State University and doctorate from Cornell.

As a weed specialist, Dr. Majek studied the growth, development, reproduction and competitive ability of weeds, using this information to develop new weed control practices. He evaluated the effectiveness of crop response to and cost potential of cultural, biological, mechanical and chemical methods of controlling weeds in vegetables and tree fruit. He also worked to determine the fate of herbicides in plants and the environment.

A long-time member of the Northeastern Weed Science Society and the Weed Science Society of America, Majek held many leadership positions in those organizations, including President. Prior to retiring in 2013, Majek was the Director of the Rutgers Agricultural Research and Extension Center in Bridgeton.

Since 1932, the New Jersey State Board of Agriculture has awarded the prestigious Citation for Distinguished Service to Agriculture to men and women who have given unselfishly of their time and talents



Secretary Fisher awards Dr. Brad Majek the Distinguished Service Award from the NJ Department of Agriculture.

to the advancement and betterment of the agricultural industry and rural life in New Jersey. This award is given to recognize and honor those individuals who have made outstanding contributions of public service to New Jersey agriculture.



Kurt W. Alstede Receives 2016 New Jersey State Horticultural Society Outstanding Fruit Grower's Award

Win Cowgill

New Jersey Agricultural Experiment Station

Kurt Alstede, was presented the New Jersey State Horticulture Societies Outstanding Fruit Grower's Award at the 71st North Jersey Fruit Meeting Held in Flemington NJ on February 24, 2016.

Located in Chester, New Jersey, Morris County Alstede Farms, LLC has evolved to a retail based business raising nearly 350 acres of tree fruits, small fruits, vegetables, and flowers. <http://alstedefarms.com/>

Kurt functions as the General Manager and is supported, by a strong management team that includes his wife of 19 years, Barbara.

Today 100% of the farm's output is sold through the on farm store, PYO, tailgate markets, and CSA.

Agritourism also compliments the farm's retail offerings. The farm currently employs 22 full-time year round staff members and seasonally up to 200 people in both the production and retail segments of the business.

Kurt has been an active volunteer fire-fighter for nearly 35 years and has served or is currently serving in several different capacities of local, county, state, and federal government. He is also engaged in

a variety of agricultural organizations and has served in church leadership roles.

He has served as past president of the Morris County Board of Agriculture, NJ State Horticulture Society Board of Directors, NJ Small Fruit Council Board of Directors,

Morris County Farm Service Agency, USDA, Past Committee Member and NJ Highlands Council Vice-Chairman. As a member of the Highlands council helped facilitate \$150,000 in grants to Rutgers Tree Fruit IPM program for Northern New Jersey.

Kurt has been an active member and supporter of the fruit and agricultural community in New Jersey.




From left to right: Craig Steely, Hector Ruiz, Kurt Alstede, Win Cowgill, and Peter Nitzsche.



Gary Mount, Terhune Orchards, Princeton, NJ with their new “Wine Barn” under construction, March 2016. The new 3,600 sq foot facility will be used for wine production, storage and hospitality. It also includes public bathrooms and staff lunchroom. <http://terhuneorchards.com/winery/>





Fresher, Sweeter, Juicier

Peaches From Circle M Farms



*Exceptional quality and flavor at attractive prices
Brought to you from the 4th generation family
farm of Santo John Maccherone*

- PLU Stickers Available Upon Request
- Low in Calories
- Nutritious and Delicious
- Exceptional Flavor



SANTO SAYS...

"Peaches from Circle M Farms just plain taste better. Their size and color are what your customers want, and their flavor is the icing on the cake.

That's because our orchards are tended with care and attention, and their fruit is left on the tree to ripen naturally until they reach their optimum sweetness. Then they are picked by hand, taste-tested by me to ensure quality, and packed gently into

25 lb. volume boxes for shipping. Single layer cartons are available upon request.

Circle M's 30 varieties of peaches and nectarines are known for their sweetness and their luscious color. They look as good as they taste.

Discriminating customers appreciate our white flesh peaches because of their lower acid levels and supremely sweet taste.

Add it all up: Peaches from Circle M Farms are fresher and sweeter, picked by hand when their time has come, and packed with care for a longer shelf life.

Yellow, white and donut peaches and yellow and white nectarines are available early July to late September."



**Call us at 856 856 207-4542 or
609 381-6465. Fax your order to
856 878-0404**



Available in 16 oz. and 64 oz. sizes

**Made from fresh New Jersey Peaches
"Peach Cider Drink, Peach Salsa, Peach
Preserves"**

**Santo John Maccherone
circlemfarmsllc@gmail.com**

**Circle M Farms
88 Sandy Ridge Road
Salem, N.J. 08079**



Horticultural News

New Jersey State Horticultural Society
176 Airport Road
Hackettstown, NJ 07840